

UNIVERSITY OF ILLINOIS  
LIBRARY

Class  
580.5

Book  
FB

Volume  
1  
cop. 2

Jan 09 20M

## CENTRAL CIRCULATION BOOKSTACKS

The person charging this material is responsible for its return to the library from which it was borrowed on or before the **Latest Date** stamped below.

**Theft, mutilation, and underlining of books are reasons for disciplinary action and may result in dismissal from the University.**


**TO RENEW CALL TELEPHONE CENTER, 333-8400**

**UNIVERSITY OF ILLINOIS LIBRARY AT URBANA-CHAMPAIGN**

APR 22 1992  
APR 21 1992

When renewing by phone, write new due date below  
previous due date.

79521 L162



FIELD COLUMBIAN MUSEUM

PUBLICATION 25

Botanical Series.

Vol. 1, No. 4.

---

CONTRIBUTION III  
TO THE  
COASTAL AND PLAIN FLORA  
OF  
YUCATAN.

BY


CHARLES FREDERICK MILLSPAUGH, M. D.

Curator, Department of Botany.



CHICAGO, U. S. A.

January, 1898.



## CONTENTS.


---

### I.

Plants Collected by Dr. Geo. F. Gaumer in 1895-96, Dr. Arthur Schott in 1864-66, and Mr. Witmer Stone in 1890.

### II.

Notes and New Species—Prof. Ludwig Radlkofer; A New *Myginda*—Dr. Theodor Loesener; and additional notes and species by the author.





## Third Contribution to the Coastal and Plain Flora of Yucatan.

CHARLES F. MILLSPAUGH, M. D.

---

Since the issue of the second of these contributions, this Museum has been so fortunate as to secure from the widow of the late Dr. Arthur Schott, his entire herbarium, which includes about nine hundred Yucatan plants collected by him in 1864-66 while engaged by the Mexican Government to make a geologic survey of the Peninsula. This collection is of great importance, it being the first really comprehensive attempt to collect the flora of this region. It is, however, to be regretted that Dr. Schott's idea of numeration should have been that of numbering his collections at home, instead of in the field, and attempting to group his species of whatever habitat under certain arbitrary numbers, before such species had been positively ascertained. This will cause conflict between many of the numbers published in these contributions and those in the herbaria of the National Museum and Kew, where part sets of these plants have been previously deposited by him. A majority of his plants are included in this contribution.

In 1890 an expedition was sent into Mexico under the auspices of the Academy of Natural Sciences of Philadelphia, and in charge of Prof. Angelo Heilprin. The work extended from the Orizaba region to the Peninsula of Yucatan, where Mr. Witmer Stone, the Ornithologist of the party, incidentally collected about three hundred plants. These were approximately determined by the late Mr. J. H. Redfield and distributed, as about ninety-five species, in the herbarium of the Academy, whence they have been kindly reassembled by Mr. Stewardson Brown, and communicated to me for study by the Academy. These species, so far as determined at this time, are also included here.

Even a limited opportunity to examine the collections at the herbarium of Columbia University convinces me that the collection made by the Hon. E. P. Johnson in Yucatan and Tabasco, in 1848,

is of considerably greater proportions than Professor Helmsley reports in the botanical volumes of the *Biologia Centrali-Americana*. Only a patient overhauling of the whole herbarium can definitely settle the question of the species in this collection. A few genera in Leguminosæ revealed several plants which I have incorporated in the following list.

Dr. Geo. F. Gaumer has continued his collections during the past year. These, so far as determined, form the basis of this contribution, and appear in the text in parenthesized numerals in the body type—the numbers of other collectors being in *italics*.

To the above collections are added, in the following list, a few hitherto unclassified numbers collected by the author in 1887, and in 1895 while a member of the Allison V. Armour expedition, and the remaining numbers of Sr. Porfirio Valdez' century of "Medicinal Plants" collected in 1895. As heretofore, the species appearing in **black-face** type are new to the region covered by these publications.

Those specialists who have kindly contributed to the knowledge of the species, are credited under the orders they have worked.

Plants Collected by Dr. Geo. F. Gaumer in 1895-96,  
Dr. Arthur Schott in 1864-66, and Mr.  
Witmer Stone in 1890.

---

FUNGI.

**Lentinus villosus** Klotzsch.  
Merida "in a garden," E. H. Thompson (947).

LICHENES.

**Ramalina calicaris farinacea** Schaer. **Mex.-gul.**  
On shrubs, Downs at Progreso, April to Dec. 1865 (*Schott, 311*).  
Powdery soredia and lateral apothecia both plentiful.

**Ramalina calicaris fraxinea** Fr.  
"Common at Progreso" (1175).

**Ramalina rigida** (Pers.) Tuck. Am. Lich., i. 22.  
Downs of Progreso, Dec. 1865 (*Schott, 311*). On trees and  
shrubs; tufts large, branches attenuate elongated, tips filiform,  
apothecia small.

ALGÆ.

**Nostoc verrucosum** Vauch. Hist. Alg., 225, *fide Prof. W. G. Farlow.*  
"Found in the aguada Chulubmay, nine miles east of Izamal,  
March" (1097).

CHARACEÆ.

**Chara gymnopus** A Br. *fide Prof. T. F. Allen.*  
"Common in the aguadas east of Izamal" (913). An inde-  
terminable sterile sub-species of this form.

MUSCI.

**Leucobryum incurvifolium** C. Muell. sp. nov.  
Species optima, ab omnibus congeneribus. Foliis rotundato-  
obtusatis brevissime hyalino-apiculatis, definite apice incurvi  
valde recedens.  
"Common in the forests at Buena Vista Xbac, September"  
(1117).

**Stereophyllum** (*Monerium*) **perpusillum** C. Muell, sp. nov.

Monoicum caespituli latissimi perpusilli taxicaulioidei pallide virides nitiduli intricati; caulis perbrevis vage ramosus, ramis pusillus tenuibus complanatus ætate albescenti-viridibus madore parum turgidulis; folia caulina minuta in axi crassiusculo pallide virente laxè horridè conferta humore regulariter patulâ plagiothecoidea, e basi angustiore ad alam unicum cellulis nonnullis parenchymaticis pellucidis majusculis ornata anguste oblongo-lanceolata breviter acuminata acutata integerrima, nervo angustissimo mediano viridi parum carinata-concava, e cellulis parvis longiusculis angustiusculis laxiusculis mollibus chlorophyllosis reticulata; perichætalia pauca parum minore apressa longius acuminata; theca in pedicello brevi rubente inclinata ætate nutans parva oblonga sub ore valde coarctata fusca, operculo breviter conico; peristomium pro capsula robustum, dentibus externis angustis lutescentibus cristatis, internis minutis ciliatis rudimentariis.

Habitato America-centralis, Yucatan, in Sylvis proxe Izamal, 1897 (340.)

Ex habitu ad *Stereophyllum Peruvianum* (Mtze.) accedens, sed hæc species foliis multo laxius reticulatis magis acuminatis basi ad alar utrinque parenchymaticis longe differt.

## FILICINÆ.\*

**Adiantum tenerum** Swz. Prod. 135. *Culantrillo.*

"A growth 3 feet high, in a cave about one mile northwest of Izamal" (1071). At the cenote of Kickil (*Schott*, 778). Quinta Encalada, Merida, March 26, 1865 (*Schott*, 92).

## ADIANTUM TRICHOLEPIS Fee.

At the ruins of Uxmal, Sept. 15, 1866 (*Schott*, 687). Cave near Calcehtok, March 11, 1890 (*Stone*, 263, in *Herb. Acad. Sci., Phila.*).

## ANEIMIA ADIANTIFOLIA Swz.

At Teocali, and on the ruins in Mayapan, Feb. 28, 1866 (*Schott*, 651).

**Aneimia bipinnata** Moore.

Cenote ya Usil, Hacienda de Mucuyché Oct. 20, 1865; and Cenote de Uayma, March, 1866 (*Schott*, 677).

## ASPIDIUM TRIFOLIATUM Swz. Syn. 43.

"Growth 3 feet high, abundant in a large cave at Buena Vista Xbac, November" (994). Cenote de Sta. Ana, Valladolid, March, 1866 (*Schott*, 782, immature specimens from same locality, 780) Cave near Calcehtok, March 11, 1890 (*Stone*, 261, in *Herb. Acad. Sci., Phila.*).

\*Determined by Mr. George E. Davenport.

**Asplenium dentatum** Linn.

At the Cenote de Telchaguillo, March 1, 1866 (*Schott, 747*).

## ASPLENIUM PUMILUM Swz.

Sisalchen.

Quinta de Obispo, Merida, Nov. 15, 1864 (*Schott, 11*).

## CHEILANTHES MICROPHYLLA Swz. Syn. Fil. 127.

On walls at Merida, Oct. 4, 1865; and at the ruins of Uxmal, Sept. 16, 1865 (*Schott, 686*). In well near Izamal, Feb. 28, 1890 (*Stone, 253*, in *Herb. Acad. Sci., Phila.*).

**Hemionitis palmata** Linn. Sp. Pl., 1535.

"Growth 8 inches high, common in the dense forests of Buena Vista Xbac" (1072).

**Phegopteris rudis.**

A single young frond, apparently of this species, was collected by Dr. Schott with his specimens of *P. tetragona*, (631) at the Cenote de Sacalum, Sept. 18, 1865.

**Phegopteris tetragona** Mett. Fil. Hort. Lips., 84.

"A growth 16 inches high, common on the walls of a cenote about 12 miles east of Izamal" (995). Dr. Gaumer sends another specimen of this species under the number (993) remarking: "Fern 2 feet high, occasionally seen in cultivation, and said to grow wild," which wild state is proven in his 995. Cenote de Sacalum, Sept. 18, 1865 (*Schott, 631*).

**Polypodium incanum** Swz. Fl. Ind. Occ., iii., 1645.

"Found growing upon the branches of *Lysiloma latisiliqua* at Buena Vista Xbac, in September, not seen again" (1110).

## POLYPODIUM LYCOPODIOIDES Linn.

Climbing trees at the cenote of Kikil, March 27, 1866 (*Schott, 768*).

## POLYPODIUM PHYLLITIDIS Linn.

At the cenote Mascabicú, March, 1866 (*Schott, 781*).

**Polypodium reptans** Swz.

At the cenote Uayma, March, 1866 (*Schott, 779*). Cave near Calcehtok, March, 11, 1890 (*Stone, 264*, in *Herb. Acad. Sci., Phila.*).

## SALVINIACEÆ.

**Salvinia auriculata** Aub. *vide Prof. Underwood.*

"Found floating upon the surface of the water of many of the aguadas near Izamal, November" (1007). Smaller than usual and sterile; probably young plants.

## LYCOPODINÆ.

## SELAGINELLA LONGISPICATA Underw. Field Col. Mus. Bot., i., 287.

Small but typical plants from Nohcacab, Nov. 1865; and Merida, Sept. 1865 (*Schott, 669*).

## CYCADACEÆ.

- Cycas circinnalis** Linn. Sp. Pl., 1658. *Teoiczoll.*  
Cultivated at Merida, March 11, 1866, not in flower (*Schott*,  
748.)

## GRAMINEÆ.\*

- Andropogon hirtiflorus** Kth. Revis. Gram., ii., 569.  
"Tekax, on old fields, 3 feet high, Sept." (1134.)
- Andropogon semiberbis** Kth. Enum., 489.  
"Two feet high, abundant on stone walls; house roofs and on  
the ancient ruins, rare in fields, about Izamal," Nov. (1037.)
- Andropogon sorghum Drummondii** Hack. de C. Monog. Phan.  
vi., 507.  
"Three feet high, in an old field at Buena Vista Xbac,"  
April (1053.)
- Antheophora elegans** Schreb. Besch. Græs., ii., 105.  
"Common in old fields near Izamal, 1 foot high." Oct. (1030).  
Merida, Sept.; and Hacienda San Rafael, Oct. 1865 (*Schott*, 549).
- Aristida nigrescens** Presl., Rel. Hænck., i., 223.  
Merida, Aug. and Oct. 1865 (*Schott*, 601, 656 in part).
- Arundo Donax** Linn. Sp. Pl., 81.  
"Common on the coast in aguadas and cenotes, often culti-  
vated. 20 feet high" (1141).
- Bouteloua Americana** (Sw.) Scribn. loc. cit. 306.  
*Aristida Americana* Sw. Obs. 41, t., f. 2; 1791 ex Kunth.  
*Sinebra repens* H. B. & K. Nov. Gen. et Sp., i. 172, t. 52; 1815.  
"Brushlands near Izamal, uncommon, 3 feet high." Oct. (1036).
- Bouteloua bromoides** Lag., Gen. et Sp. Nov., 5.  
Merida, Sept., Oct. and Nov., 1865; and at the ruins of  
Uxmal Sept. 16, 1865 (*Schott*, 656, 742, 659). Depauperate  
specimens from Tekanto, Feb. 28, 1890 (*Stone*, 189, in Herb.  
Acad. Sci. Phila.)
- Bouteloua disticha** (H. B. K.) Benth. Jour. Linn. Soc. xix., 105.  
*Polyodon distichum* H. B. K. Nohcacab, Nov. 24, 1865 (*Schott*,  
741). Presl. referred this to *Bouteloua bromoides* H. B. K.
- BOUTELOUA TRIÆNA** (Spreng.) Scribn. Proc. Phila. Acad. 1891, 307.  
*Triæna racemosa* H. B. K. "Two feet high; uncommon on  
old fences and low shrubbery, upon which it depends for sup-  
port." Izamal, Aug. (1026). Nohcacab, Nov. 24, 1865 (*Schott*,  
740) Tekanto, Feb. 25, 1890 (*Stone*, 210, in Herb. Acad. Sci.  
Phila.).  
For synonymy see p. 10, Contrib. I.

\*Determined by Prof. F. Lamson-Scribner.

**Cenchrus echinatus** Linn. Sp. Pl., 1050.

To be compared with *C. viridis* Spreng., which Griesebach refers to *C. echinatus* as a variety. "Very abundant on cultivated lands near Izamal, 18 inches high" (1084). Merida, Aug. 1865 (*Schott, 498* in part).

**CENCHRUS PALLIDUS** Fourn., Mex. Pl. Enum. Gram., 50. **Mul.**

Merida, July and Aug. 1865 (*Schott, 596, 498*). *Schott's 596* equals Liebmann's 465.

**Chloris barbata** Sw. Fl. Ind. Occ., i. 200.

"Abundant on stone walls near Izamal, 18 inches high." (1085).

**Chloris ciliata** Sw., Prod. Veg. Ind. Occ., 25.

Merida, Aug. 3, 1865 (*Schott, 383*). Calcehtok, Mar. 11, 1890 (*Stone, 274*, in Herb. Acad. Sci. Phila.).

**Chloris elegans** H. B. K., Nov. Gen. et Sp. i. 166.

At Sacalum, Sept. 18, 1865 (*Schott, 632*).

**COIX LACHRYMA-JOBI** Linn. Sp. Pl. 972.

"Izamal. One plant came up spontaneously in my yard in 1894, and produced a quantity of seed; this was followed in 1895 by several plants springing up in the same place and producing a great quantity of seed; but none came up in 1896" (1031). See p. 9 spec. cit.

**CYNODON DACTYLON** Pers., Syn. Pl., i. 85.

At the hacienda Saragossa, Sept. 8, 1865 (*Schott, 737*).

**Dactyloctenium Ægyptiacum** (L.).

Merida, Aug. and Sept. 1865 (*Schott, 734*). *Cynosurus Ægyptius* Linn.

**Distichlis spicata** (L.). Greene Bull. Calif. Acad., ii. 415.

*Uniola spicata* Linn. Celestun, May 13, 1865 (*Schott, 494*).

**ELEUSINE INDICA** Gaertn. Fruct., i. 8.

"Abundant in shady places near Izamal, 15 inches high" (1086). Merida, Aug. and Sept. 1865 (*Schott, 593* in part, 738).

**ERAGROSTIS CILIARIS** Link. Hort. Berol., i, 192.

Yucatan loc. ignot. (851). Merida, at Quinta del Obispo, Nov. 24, 1864, and Sept. 5, 1865; and at the ruins of Uxmal, Sept. 16, 1865 (*Schott, 20, 619*). Tekanto. Feb. 27, 1890 (*Stone, 187*, in Herb. Acad. Sci. Phila.).

**Eragrostis elongata** Jacq. Ec. Gram., t. 3.

Sublitoral at Sisal, Oct. 24, 1865 (*Schott, 640*). Compares with specimens in the Bernhardt herbarium, labeled by Presl.

**Eragrostis plumosa** Link, Hort. Berol., i. 192).

Merida at Quinta del Obispo, Nov. 1864, and Sept. 1865 (*Schott, 59*).

**Guadua latifolia** H. B. K.? Syn. Pl. i. 254.

Merida, Jan. 1866, lacking inflorescence (*Schott, num. amiss.*).

**Gouinia latifolia** (Griseb.) Vasey. Rose in Contrib. Natl. Herb., i. 365.

*Tricuspis* (Neuroblepharum) *latifolium* Griseb., Pl. Lorentz, 259. Nohcacab, Nov. 25, 1865 (*Schott, 703*).

**Gouinia virgata** (Presl.) Scribn. U. S. Dept. Agr. Div. Agrost. Bull. 4, 10.

*Bromus virgatus* Presl. Rel. Hænk., 263. *Gouinia polygama*. Fourn. Enum. Pl. Mex. Gram., 103. Tekax, Sept. (1033).

**Ichnanthus lanceolatus** Scribn. & Sm.

**Xcanchim.**

“An erect or ascending, cæspitose, branching perennial 1 to 2 feet high, with lanceolate leaves and simple panicles of few loosely flowered racemes. Sheaths shorter than the internodes, ciliate along the margins, otherwise smooth, or the lowermost pubescent; ligule a short ciliate fringe of hairs; leaf blade 1 to 3 inches long,  $\frac{1}{4}$  to  $\frac{1}{2}$  inch wide, lanceolate acute, smooth, many nerved, with a narrow cartilaginous margin abruptly narrowed at the base, this contraction forming in the lower leaves, especially those of the sterile shoots, a slender channeled petiole, which, like the sheaths, is ciliate along the margins. Panicle branches, erect or ascending (spreading in anthesis), 1 to 2 inches long, the uppermost shorter. Spikelets in pairs, one subsessile, the other raised on a pedicel about as long as itself. Spikelets ovate-lanceolate, acute, glabrous, about 2 lines long; first glume ovate, acute, strongly 3-nerved, one-half to three-fourths the length of the spikelet, scabrous on the mid-nerve above; second glume ovate-lanceolate, acuminate, 5-nerved, nearly clasping the similar empty third glume; base of the fourth glume surrounded by the third; fourth glume about 1.5 lines long, oblong-lanceolate obtuse, 5-nerved, very smooth and closely rolled about the palea, which is of similar texture.—“Old fields about Izamal, No. 854. George F. Gaumer, September, 1895. Yucatan.” Scribner & Smith, U. S. Dept. Agric. Div. Agrost. Bull. 4, Feb. 6, 1897, p. 36. “Common in the forests of Buena Vista Xbac,” Sept. (1111). This is the *Panicum* sp. of p. 288, these contributions.

**LEPTOCHLOA MUCRONATA** Kth. Rev. Gram. i, 91.

Environs of Merida (*Schott, May 1865, 401; July 1865, 401, 590; Nov. 1865, 739.*)

**Oplismenus Burmannii** (R. Br.) Beauv., Agrost., 54.

*Orthopogon* R. Br., Prod., 194. “In shady places, 3 feet high, common about the mouths of caves and around cenotes and aguadas near Izamal,” Oct. (1038). Merida at the Quinta del Obispo, Nov. 1864; and Merida, July 1865 (*Schott, 55, 384.*)

In Dr. Schott's 384 and Dr. Gaumer's 1038 the spikelets are somewhat larger and the foliage larger and denser than in Dr.



Schott's 55, and some authors have referred the former to *Oplismenus cristatus* (Presl). Both forms have been referred to *O. Humboldtianus* Nees. In the Kew Index, *O. cristatus* and *O. Humboldtianus* are referred to *O. Burmannii*. Beauv.

**Panicum Carthaginense** Sw., Fl. Ind. Occ., i. 148.

Merida, July 11, 1865 (*Schott*, 592).

**Panicum colonum** Linn., Syst. ed. x, 870.

At the Quinta del Obispo, Merida, Nov. 1864; and at Merida, June and July 1865 (*Schott*, 42, 598).

**Panicum Crus-galli** Linn. Sp., Pl., 50.

Borders of an aguada at Kanachen, March 1866 (*Schott*, 832).

**Panicum divaricatum** Linn. Elmg. Pl. Jam. Perg., 9.

"Common on fences in low brushlands, 4 feet high. Izamal" (1032). Merida, 8 to 9 feet high, Oct. 11, 1865 (*Schott*, 675).

**Panicum divaricatum latifolium** (L.) Fourn. Enum. Pl. Mex. Gram., 33. Sit.

*P. lanatum* Sw. Griseb., Flor. Brit. W. Ind., 551. "Common along old fences and in brushlands near Izamal, Aug. to Feby." (1025.) Merida, Aug. and Sept. 1865 (*Schott*, 600).

**Panicum fuscum** Sw., Prod. Veg. Ind. Occ., 23.

Merida, May and June 1865 (*Schott*, 384).

**Panicum lanatum** Rottb. Act. Lit. Univ. Hafn., i. 269.

*P. insulare* Meyer., *P. leucophæum* H. B. K. Merida, 1865 (*Schott*, 94).

**Panicum marginatum** (Link) not R. Br., nor Vahl.

*Digitaria marginata* Link. Hort. Berol., i, 102. *Panicum Linkianum* Kth. Gram., i. 33. "Common in waste places near Izamal. 3 feet high, Nov." (1034.) Merida, Aug. 27, 1865 (*Schott*, 609).

**Panicum maximum** Jacq. Ic. Pl. Rar., i, t. 13. *Sacate Guinea*.

*P. jumentorum*. Pers. Syn., 83. "Common in cultivation, growing from 6 to 8 feet high. Izamal, June" (719). Merida, Aug. 20, 1865 (*Schott*, 595).

**Panicum sanguinale** Linn. sp. Pl. 57.

Yucatan, loc. ignot. (1029). Merida Aug. 20; ruins of Uxmal, Sept. 16, 1865 (*Schott*, 586, 735).

**Paspalum elongatum** Griseb. Pl. Lorentz., 260.

Merida, Aug. 3, 1865 (*Schott*, 593). This specimen agrees with 1306 Bang's Bolivian collection of May 1892.

**Paspalum lentiginosum** Presl.? Rel. Haenk., i. 218.

Merida, July 11, 1865 (*Schott*, 597). Allied to *P. paniculatum* Linn. Spikes 4 to 5 slender, 1.5 to 2.5 in. long, the lower ones remote.

**Paspalum paspaloides** Pers. Syn., 81.

"Izamal, common in old fields, 3 feet high, Oct." (1027).

**Paspalum vaginatum** Sw., Prod. Veg. Ind. Occ., 21.

In fresh water at the aguada and Cenote de Nabalá on the Hacienda Chablé, Aug.; and salt swamps near Sisal, Nov. 1865 (Schott, 546, 733).

This species appears to me distinct from *P. distichum* Linn.

## SETARIOPSIS AURICULATA (Fourn) Scribn., Field Col. Mus. Bot., 1. 289.

Merida, July and Aug.; and Hacienda Saragossa, Sept. 1865 (Schott, 592, 591, 621).

**Streptachne tenuis** H. B. & K., Nov. Gen. et sp., 1, 124.

*Aristida tenuis* Kth. Rev. Gram., 1, 62. "Uncommon among low shrubbery along old fences near Izamal, 3 feet high. Aug." (1024.) Merida, Sept. 1865 (Schott, 736).

Under the number 594 Dr. Schott collected in the Aguada de Labcah in July 1865, a portion of a gramineous species near *Arundinaria*, to which he applies the Yucatec name *Cañote*, also 547 from gardens at San Rafæl de Xteppen, an *Andropogon*. Neither specimen bears flowers nor fruits, in fact no portion of an inflorescence. With these two the above list includes all the gramineæ in the Yucatan portion of his herbarium.

## CYPERACEÆ\*.

**Cyperus** Sp.

Prox. *speciosus*. "Height 3 feet, common in moist, shady places near Izamal." (1035).

## CYPERUS OCHRACEUS Vahl. Enum. ii., 325.

Yucatan loc. ign. (1028).

**Cyperus rotundus** Linn. sp. Pl., 67?

"One foot high, common in moist places near Merida" (1088).

**Cyperus squarrosus** Linn. Cent. Pl., ii. 6.

"Six to 10 inches high, common in old fields near Izamal" (1023).

**Rynchospora micrantha** Val. Enum. ii., 231.

"One foot high, common in the forests of Buena Vista Xbac, September" (1115).

## PALMÆ.

**Acrocomia Mexicana** Karw. Mart. Hist. Nat. Palm, iii. 285 *Cocoyol*.

Merida, Dec. 1865 (Schott, 805). "Next to the Coco the tallest and most graceful palm of the region."

\*Determined by Prof. N. L. Britton.

**Attalea Cohune** Mart. in Palmet. Orb., 121. *Caraso.*  
Cult? Hacienda Oxcom, Feb'y 4, 1866 (*Schott*, 726 in Herb.  
U. S. Nat'l Mus.).

**Chamædorea gracilis** Willd. Sp. Pl. iv. 800. *Xiat.*  
"Palm 20 feet high, abundant at Xcholac and Buena Vista  
Xbac, flowering from Feb'y to March" (420). Ruins of Nohpat,  
Nov. 24, 1865 (*Schott*, 719).

**COCOS NUCIFERA** Linn. Sp. Pl., 1658.  
Cult. Merida, Dec. 1865 (*Schott*, 712).

**Phoenix dactylifera** Linn. Sp. Pl., 1658.  
Cult? at the Rancho San Joaquin, near Mayapan, March 1,  
1866 (*Schott*, 749).

**SABAL MEXICANUM** Mart Hist. Nat. Palm., iii 246. *Xaan.*  
"A very common tree in the forests of the peninsula, flower-  
ing throughout the year" (317). Merida, Jan'y, 1866 (*Schott*,  
802).

**Thrinax argentea** Lodd. Desf Cat. Hort. Par., iii. 31. *Chit.*  
Downs at Progreso, Dec. 22, 1865 (*Schott*, 293, 721).

## ARACEÆ.

**Arisæma** sp.  
"In moist places, forests about Izamal" (1091).

**Arum Italicum** Mill. Gard. Dict., ed. viii, n. 2.  
Merida Cult? March to June, 1865-6 (*Schott*, 489).

**Caladium bicolor** Vent. Jard. Cels., t. 30. *Papagayo.*  
Merida, June, 1865 (*Schott*, 434).

**Philodendron trifoliatum.**  
Mayapan, Feb'y 28, 1866 (*Schott*, 836).

## LEMNACEÆ.

**WOLFFIA BRAZILIENSIS** Wedd. An. Sc. Nat., 1849. 170.  
Hacienda de San Rafael Xteppen, July 7, 1865, and Hacienda  
de Chablé, Aug. 8, 1865 (*Schott*, 541).

## BROMELIACEÆ.

**ÆCHMEA BRACTEATA** (Sw.) Mez. de C. Monogr., ix. *Xkeo.*  
"Six feet tall, abundant in the forests about Izamal, producing  
its yellow flowers from Feb'y to June" (419). Road to Progreso,  
April 7, 1865 (*Schott*, 274).

**BROMELIA PINGVIN** Linn. Sp. Pl., 285. *Oalbay.*  
Aguada Calatcapek, April 2, 1866 (*Schott*, sine. num.).

**Karatas Plumieri** E. Morr. Belg. Hort. 1872, 131. *Pinuela*. **Chom.**  
Merida, June 19, 1865 (*Schott, 558*).

**Tillandsia brachycaulos** Schl. in *Linnaea*, xviii, 422. **Mis.**  
“Herb 12 inches high, abundant on trees near Izamal, producing its royal-purple flowers in June” (742). Merida, July, 1865 (*Schott, 557*).

TILLANDSIA FASCICULATA LATISPICA Mez. de C. Monogr. Phan, 9.  
Merida, March 10, 1865 (*Schott, 435*). **Xolohbenal.**

TILLANDSIA RECURVATA Linn. Sp. Pl., ii, 410.  
Merida, Nov. 13, 1864 (*Schott, 35*).

**Tillandsia setacea** Sw. Fl. Ind. Occ. i, 593.  
Merida, July 11, 1865 (*Schott, 842*).

TILLANDSIA STREPTOPHYLLA Scheidw. Hort. Belge. iii, 252.  
Merida, June 26, 1865 (*Schott, 349*).

TILLANDSIA UTRICULATA Linn. Sp. Pl., ed. ii, 411.  
Merida, July 28, 1865 (*Schott, 556*).

TILLANDSIA VESTITA Cham. & Schl. *Linnaea* vi, 52.  
Merida, Dec. 11, 1864 (*Schott, 161*).

#### COMMELINACEÆ.

CALLISIA REPENS Linn. Sp. Pl., 62.  
Merida (*Schott sine. num.*) ; loc. ign. (*Schott, 151*). “Herb 18 inches to 2 feet, abundant in fields and forests about Izamal, its blue flowers appearing throughout the year” (376). “Prostrate 3 feet, rooting at the joints (1079).

**Commelina nudiflora** Linn. Sp. Pl., 4.  
Merida. Nov. 30, 1864 (*Schott, 32*).

**Commelina pallida** Willd. Hort. Berol. ii, 87.  
Collected by Dr. Gaumer near Izamal together with *Callisia repens* No. 376.

RHOEO DISCOLOR (L. Her.) Hance, Walp. Ann., iii, 659. **Chac-Oam.**  
Merida, Feb'y 11, 1865; Ruins of Labcah near Hacienda Chemax, July 30, 1865 (*Schott, 239, 564*).

**Spironema fragrans** Lindl. Act. Soc. Sc. Fenn, x, 127.  
“One foot high, rare at Xcholoc, producing its white flowers in March” (421).

**Zebrina pendula** Schniz. in Bot. Zeit vii, 870.  
“Ascending 3 feet, commonly cultivated at Izamal, producing its bright blue flowers from August to March” (412).

## LILIACEÆ.

**Smilax Gaumerii** sp. nov.**Xcoché.**

Stem cylindrical, striate, climbing 25 feet, armed with strong triangular-flattened prickles, mostly in pairs opposite the branches, sometimes slightly curved at the points; branches sub-cylindrical, striate, more or less zig-zag, aculeate; leaves ovate-lanceolate, taper-pointed, mucronate, oblique at the base, strongly reticulate-veined above and below, 3-nerved, the mid rib furnished with 1-2 recurved prickles; petioles short, jointed at the middle and furnished with a stout recurved prickle at the joint; stipules sheathing, often aculeate and sometimes prolonged into light grey ligneous tendrils. Inflorescence axillary, globular, peduncle about half the length of the pedicels. Leaves 1 to 2.5 inches long, .5 to 1 inch broad, petioles .25 inch long, stem prickles .25 to .36 inch long. Frequent on forest and brush lands about Izamal, June (687).

## AMARYLLIDACEÆ.

**YUCCA YUCATANA** Engelm Trans. Acad. St. Louis iii. 37.

Ruins of Nohpat, Nov. 24, 1865 (*Schott*, 706).

**Zephyranthes Lindleyana** Herb. Amaryll. 174. t. 35. f. 5.

"Herb 8 inches high, very abundant in the town of Izamal, opening its sulphur-yellow flowers in September (836).

## DIOSCOREACEÆ.

**Dioscorea alata** Linn. Sp. Pl., 1033.

Hacienda San Rafael Xteppen, Cult?, Aug. 7, 1865 (*Schott*, 548).

**Dioscorea calyculata** Donnell-Smith, Bot. Gaz. xx, 295.

"Vine 40 feet, common on vacant and waste lands where it flowers in November" (931).

**Dioscorea floribunda** Mart. & Gal., Bull. Acad. Brux., ix, 391.

"Vine 20 feet, common on brush lands, Nov." (925, 926).

**Dioscorea macrostachya** Bth., Pl. Hartw., 73.

"Vine 12 feet, abundant on brush lands, Sept." (794).

**Dioscorea polygoniodes** H. & B., Willd., Sp. Pl., 795.

"Vine 20 feet, common on brush lands, Nov." (928).

**Dioscorea sativa** Linn. Sp. Pl., 1033.

**Bolador.**

"Vine 60 feet, rarely cultivated; blooms in October, and produces at every leaf an edible fruit resembling the potato, the large tuberous root is also eaten" (920).

## MUSACEÆ.

- MUSA SAPIENTUM** Linn. Sp. Pl. 1477. **Haas.**  
Merida at the Quinta del Obispo July to September, 1864-5  
(Schott, 849).

## CANNACEÆ.

- Canna edulis** Ker-Gawl Bot. Reg. ix, t. 775.  
"Common in cultivation at Izamal where it flowers from July  
to March" (963).

## PIPERACEÆ.

- Piper auritum** H. B. K., Nov. Gen. & Sp., i. 54. **Xmacolan.**  
Merida at the Quinta del Obispo, Nov. 25, 1864 (Schott, 183).  
Abundant about Merida (Valdez, 80).

## CASUARINACEÆ.

- CASUARINA EQUISETIFOLIA** Linn. Amoen. Acan., iv. 143. *Cipres.*  
A cultivated tree 100 feet high, loc. ign. (945). Merida Dec.  
1864 (Schott, 60).

## MORACEÆ.

- ALICASTRUM BROWNEI** Kuntze, Rev. Gen., 623.  
*Brosimum Alicastrum* Sw. Merida, June 25, 1865; Hacienda  
Xteppen, July 7, 1865 (Schott, 555, 561). Tekanto, Mar. 9, 1890  
(Stone, 225, in Herb. Acad. Sci., Phila.).

- Artocarpus incisa** Linn. f. Suppl., 411. *Arbol de Pan.*  
Cult? Hacienda Taulum, Dec. 1865. (Schott, 816).

- Castilloa elastica** Cerv. Supl. Gaz. Lit. Mex., 7. *Hule.*  
Cult? Hacienda de Kanachen, March 1866 (Schott, 777).

- Cecropia obtusa** Trec. Ann. Sc. Nat., 1847, 79. **Xcoochlé.**  
Ruins of Uxmal, Sept. 15, 1865 (Schott, 639).

- Chlorophora tinctoria** (L) Gaud. Freyc. Voy. Bot. 1826.  
*Morus tinctoria* L., *Maclura tinctoria* Don. Male tree at  
Merida July 3, 1865 (Schott 517); female at the Aguada de  
Labcah, near the Hacienda de Chemax, July 30, 1865 (Schott,  
607).

- DORSTENIA CONTRAJERVA HOUSTONIANA** Bureau. **Xcampahuy.**  
Merida at Quinta del Obispo, Nov. 24, 1864 (Schott, 46).

- Ficus** sp. **Saccabah.**  
Aguada de Calotyaxek, leaves only, March 1866 (Schott, 783).

- Ficus Bonplandiana** Miq. An. Mus. Bot. Lugd., iii, 298.  
*Golondrino. Alamo.*  
Cult? at Celestun, May 12, 1865 (Schott, 352).

**FICUS LONGIPES** Miq. loc. cit.

*Copó. Alamo.*

At the Hacienda de San Rafael Xteppen Aug. 7, 1865 (*Schott, 846*).

**MORUS CELTIDIFOLIA** H. B. K. Nov. Gen. et sp., ii, 33.

Merida (*Schott, 847*).

#### URTICACEÆ.

**Urera baccifera** (L.) Gaud. in Freyc. Voy. Bot., 497. **Laal Jimin.**

*Ortega de Caballo.*

“Common on the ancient mounds, and frequently cultivated for hedges near Izamal, November, but often out of season” (936). Our plant agrees with Regnell’s 410, Minas Geraes, Brazil; and Rusby’s 1467, Mapiri, Bolivia.

**URERA MICROCARPA** Wedd Arch. Mus. Par., ix, 156.

**Laal.**

Tizimin, March 28, 1866 (*Schott, 796*).  
*Ortega de los Caballos.*

#### ARISTOLOCHIACEÆ.

**ARISTOLOCHIA BREVIPES**, Benth. pl. Hartw., 15.

*Guaco.*

Merida, Nov. 29, 1864; Aug. 30, 1865 (*Schott, 426*). Citas  
March 31, 1866 (*Schott, 839*).

**Aristolochia maxima** Jacq. Enum. Pl. Carib. 30. *Guaco del Sur.*

Nohcacab, Sept. 1865 (*Schott, 789*).

#### POLYGONACEÆ.

**ANTIGONON LEPTOPUS** Hook. & Arn. Bot. Beech. Voy., 308.

*Flor de San Diego.*

“Vine 60 feet, common in cultivation, often also found growing wild, flowers profuse and showy” (1132). Over old fences about Merida, Apr. 14, 1887 (*Millspaugh, 43*). Merida, Nov. 1865 (*Schott, 51*).

**PODOPTERUS MEXICANUS** H. & B. Pl. Æquin. ii, 89.

**Saciba.**

Merida June 16, 1865 (*Schott, 487*).

#### CHENOPODIACEÆ.

**ATRIPLEX CRISTATA** Humb. & Bonpl. ? ex Willd. Sp. Pl. iv 959.

“Common on the beach at Progreso” (1169). Our plant does not exhibit the serrations of the leaf given in the description, being sub-entire.

**Chenopodium album** Linn. Sp. Pl., 319.

*Quilites.*

“Herb 6 to 10 feet high, often observed in cultivation and escaped into fields where, however, it finally dies out” (1065).

**Salicornia Bigelovii** Torr. Bot. Mex. Bd., 184.

*S. mucronata* Bigel. non Lag. "Shores of lagoon at the Port of Silam" (633). Celestun, May 12, 1865 (*Schott, 297*).

**Suaeda fruticosa** Forsk. Fl. Æg. Arab. 70.

"Herb 2 feet high, common at Progreso" (1172).

#### AMARANTACEÆ.

**Alternanthera obovata** (Mart. & Gal.)

"Herb 2 feet high, rare, only one clump seen, and that in the Plaza de los Remedios at Izamal, where it flowers from September to January" (905).

**Alternanthera repens** (L.) Kuntze Rev. gen. Pl., 540.

"Prostrate herb 2 to 8 inches, infrequent, roadside six miles east of Izamal, April" (1070).

ALTERNANTHERA STRAMINEA (Mart.) Millsp. Field Col. Mus. Bot. i, 16.

Calcehtoh, March 11, 1890 (*Stone, 266* in Herb. Acad. Sci., Phila.).

AMARANTUS HYBRIDUS Linn. Sp. Pl., 1406.

"Herb 8 to 12 feet high, common in old fields about Izamal, flowering in December" (1081). The densely flowered, congested-paniculate form. Merida, July 1, 1865 (*Schott, 817*). Calcehtoh, March 11, 1890 (*Stone, 262*, in Herb. Acad. Sci., Phila.).

GOMPHRENA DECUMBENS Jacq. Hort. Schœnber., t. 482.

Merida, Aug. 20, 1865 and Nov. 15, 1864 (*Schott, 82, 818*)  
Tekanto, Feb. 27, 1890 (*Stone, 221*, in Herb. Acad. Sci., Phila.).

**Gomphrena globosa** Linn. Sp. Pl., 224.

The white-flowered form Merida Jany. 14, 1866 (*Schott, 731*); the purple-flowered form from the same locality (*Schott, 730*).

IRISINE CELOSIODES Linn. Sp. Pl., 1456.

"Climber 15 feet, abundant in open wild lands, near Izamal, and occasionally seen in cultivation, April" (1078).

**Kokera paniculata** (L.) Kuntze Rev. Gen. Pl., 542.

*Chamissoa* H. B. K. "A scandent shrubby plant 10 feet high, common near the aguada de Xcholac, flowering in November" (1011).

PHILOXERUS VERMICULATUS (L.) R. Br. Prod., 416.

Celestun May 13, 1865 (*Schott, 380*).

#### PHYTOLACCACEÆ.

PETIVERIA ALLIACEA Linn. Sp. Pl., 486.

Suburbs of Merida, Nov. 14, 1864 (*Schott, 31*).

*Zorilla.*



## NYCTAGINACEÆ.

ALLIONIA VIOLACEA Linn. Syst., ed. x, 890.

Hacienda Sozil northeast of Merida July 30, 1865 (*Schott*, 531).

BOERHAAVIA ERECTA Linn. Sp. Pl., 4.

"Erect, flowers lilac, Merida June 12, 1865 (*Schott*, 478) Tekanto, flowers pink, Feb. 27, 1890" (*Stone*, 214, in Herb. Acad. Sci., Phila.).

**Bœrhaavia scandens** Linn, Sp. Pl., 4.

"Procument, flowers yellow, calyx glandular, Merida, June 12, 1865" (*Schott*, 21).

MIRABILIS JALAPA Linn. Sp. Pl., 252.

Campo, suburbs of Uman, April 1887 (*Millspaugh*, 36).

PISONIA ACULEATA Linn. Sp. Pl., 1511.

Merida suburbs. In flower Dec. 1864; in fruit Feby. 1865 (*Schott*, 143). Ticul, bush 6 ft., flowers greenish-yellow, fragrant, March 16, 1890 (*Stone*, 285, in Herb. Acad. Sci., Phila.).

## PORTULACACEÆ.

PORTULACA OLERACEA Linn. Sp. Pl. 638.

"Abundant on cultivated grounds, producing its yellow flowers in profusion from July to February" (1001).

PORTULACA PILOSA Linn. Sp. Pl. 639.

Merida suburbs, Dec. 1864, July 1865 (*Schott*, 150).

SESUVIUM PORTULACASTRUM Linn. Sp. Pl., 684.

**Oaicán.**

Celestun May 1865 (*Schott*, 488).

## NYMPHÆACEÆ.

CASTALIA AMPLA (De C.) Salisb. Parad. Lond., 1805. 73.

Aguada de Labcat near Merida July 30, 1865, and Aguada de Xcholac, March 1865 (*Schott*, 528).

## ANONACEÆ.

ANONA MURICATA Linn. Sp. Pl., 536.

*Guanabano.*

Cult? Quinta del Obispo, Merida, (*Schott*, *sine num*).

**Anona palustris** Linn. Sp. Pl., ed. ii., 757.

*Corcho. Mag.*

Aguada Calotyaxek, *sine die* (*Schott*, 19).

ANONA SQUAMOSA Linn. Sp. Pl., 537.

*Saramuyo.*

Merida, cult? June 18, 1865 (*Schott*, 497), Hacienda Taulum, Dec. 1865, not in flower (*Schott*, 820).

## MENISPERMACEÆ.

CISSAMPELOS PAREIRA Linn. Sp. Pl., 1473.

Campo about Merida, in full flower June to Nov. 1864-5 (*Schott*, 91, 500).

## LAURACEÆ.

CASSYTHA FILIFORMIS Linn. Sp. Pl., 530.

Climbing over *Conacarpus erectus* at Sisal Nov. 10, 1865 (*Schott*, 698).

PERSEA GRATISSIMA Gaertn. Fruct., ii. 222.

Cult ? Merida, Feb. 6, 1865 (*Schott*, 203).

On.

## PAPAVERACEÆ.

ARGEMONE MEXICANA OCHROLEUCA Lindl. Bot. Reg., t. 1343.

Suburbs of Merida, Jan. 27, 1865 (*Schott*, 198).

## CRUCIFERÆ.

CAKILE MARITIMA Scop. Fl. Carn., ed. 2. ii, 35.

*C. aequalis* L'Her. Celestun, May 13, 1865 (*Schott*, 376).

**Raphanus Raphanistrum** Linn. Sp. Pl. 669.

"Eighteen inches high. A large number of plants came up in the plaza of Izamal in February, 1896, just after Orin's circus troupe had camped a week on the same place, and as they fed American and Mexican hay to their animals it is probable that the seed of this plant was introduced by them. Later the plant was taken under similar circumstances in Merida" (1064).

## CAPPARIDACEÆ.

**Capparis Cynocephalophora** Linn. Sp. Pl., ed. ii, 721.

Xbayunak.

Merida, May 6, 1865 (*Schott*, 351).

CLEOME SPINOSA Jacq. Enum. Pl. Carib., 26.

Merida, April 13, 1865 (*Schott*, 329).

CRATEVA GYNANDRA Linn. Sp. Pl. ed., ii, 636.

Merida (Cult ?), Dec. 1864; Feb'y 1865 (*Schott*, 156).

## MORINGACEÆ.

MORINGA APTERA Gaert. Fruct., ii, 315.

Cult. Merida at Quinta del Obispo, March 1, 1865 (*Schott*, 471).

## CRASSULACEÆ.

BRYOPHYLLUM PINNATUM (Lam.) S. Kurz. Jour. Assoc. Beng., xi. 52.

Cenote of Tunkax, March 26, 1866 (*Schott*, 770).

## LEGUMINOSÆ.

ACACIA FARNESIANA (L.) Willd. Sp. Pl. iv. Par., ii, 1083.

"A common tree 30 feet high in the forests about Izamal, producing its fragrant orange-yellow flowers in October." (1010.)  
Our plant agrees with Schott's (118) Cuba (*A Cavenia* Bert.).

ACACIA SPADICIGERA Ch. & Sch., in Linnæa v. 594.

Subin.

Merida, June and Sept., 1865 (*Schott*, 392). Xcholac common, bush 6 ft., Mar. 2, 1890. (*Stone*, 248, in *Herb. Acad. Sci. Phila.*)

ÆSCHYNOMENE AMERICANA Linn. Sp. Pl., 1061.

Yucatan loc. ignot. (*Johnson* 45<sup>b</sup>). Yucatan loc. ignot. (*Schott*, 863).

**Æschynomene Americana depila** var nov.

Characters of the species, except that the lomentis instead of being setose-hispid (*asperis*. *Hirsutis*. Sloane) are perfectly smooth.

"Herb 2 feet high, abundant in damp places along roads and in open lands near Izamal Oct." (955).

To this variety the following plants are referred: *Palmer* Aca-pulco 102 Manzanillo 901; *Heyde* 311, *Heyde & Lux* 6103 Guatemala; *Sintenis* 5560 Porto Rico; *Hitchcock* Port Marant, Constant Spr., Lucea, Jamaica; *Lloyd* Jamaica 1050; *Wilson* Jamaica 173. *Rovirrosa* 384 Tabasco; *Botteri* 363 Orizaba, *Skinner*, Guatemala; *Wright* 124 or 125 Cuba; *Bernardi* *Hb.* San Domingo 184; *Anderson* Trinidad; *Hayes* Panama 767; *Eggers* 202 St. Thomas; *Fendler* 1779 Venezuela; *Ricksecker* 133 St. Croix; and *Donnell-Smith* 2283 Esquintla. In the following specimens Mr. Greenman notes that although the lomentis are more or less setose-hispid in the early stages they later become essentially glabrous; *Sintenis* 374<sup>b</sup> Porto Rico; *Pringle* 2515 Jalisco, *Donnell-Smith* 4160 Chupadero.

Dr. Rose in his enumeration of Dr. Palmer's 1890-91 Mexican collection (*Contrib. U. S. Natl. Herb.*, 1. 321) calls attention to this form, *Palmer* 901 Manzanillo.

ÆSCHYNOMENE FASCICULARIS Cham. & Schl. in Linnæa, v. 584.

"Ruins of Uxmal Sept. 16, 1865." (*Schott*, 864 in *Herb. U. S. Natl. Mus.*) Merida June 11, 1865, and on road from Merida to Progreso May 7, 1865 (*Schott*, 276). "Shrubby, 3 feet high, common on roadsides near Tekax, in blossom September" (1126).

**Amerimnon Brownei** Jacq. Pl. Carib., 1760.

*Dalbergia Amerinum* Benth. "A rare shrub in the forests about Izamal, its reddish-purple flowers appearing in June." (728).

**Arachis hypogæa** Linn. Sp. Pl., 741.

Peanut.

"Herb 2 to 3 feet high, common about old farm lands near Izamal, where it blossoms in October" (997).

**Bauhinia Cavanillei** nom. nov.**Chacouludtok.**

*Pauletia inermis* Cav. Ic. t. 409, 1799; *Bauhinia inermis* (Cav.) Pers. Syn. i, 455, 1805 non Forsk. Fl. Ægypt, Arab. 85, 1775.

Nohcacab and vicinity, Nov., 1865, leaves sudorific, wood very flexible; Kabah, Nov. 24, 1865 (*Schott, 702*).

**BAUHINIA LATIFOLIA** Cav. Ic. v. 4, t. 405.

Tunkas, 4 feet high, flowers white; Ticul, 6 feet high, March 3 and 16, 1890 (*Stone, 228, 246 and 284*, in Herb. Acad. Sci., Phila.).

**BAUHINIA PORRECTA** Sw. Prod. Veg. Ind. Occ., 66.

Merida suburbs and Quinta Encalada, Nov. 29, 1864, and March 26, 1865 (*Schott, 8*).

**Bauhinia splendens** H. B. K., Nov. Gen. et Sp. vi, 319.

Yucatan loc. ignot., 10 to 15 feet high (*Johnson, 32, visa* in Herb. Torrey).

**Bradburya Schottii** sp. nov.

Caule scandente glabriusculo, foliis longe petiolatis 3-foliolatis, foliolis hastatis vel subhastatis mucronatis margine pilosis, stipellis geminiis linearis, pedicellis foliis æquantes solitariis 1-floris, bracteis ovato-lanceolatis calyce sæpe, longioribus calycis cupuliformis sub 5-fidis: lobis inferioris majusculis, superioris coalitis, lateralis triangularis. Leguminibus sessilis, longe apiculatis, costæ prominente percursis inter margine et pars media, semine compressa. Caule 10 pedales, foliis et pedicellis 5 poll. long., foliolis 2.5 poll. long., 2 poll. lat., flores 1.5 poll. long., leguminis 5 to 6 poll. long., 3 lin. lat. Ad Nohpat, Nov. 24, 1865 (*Schott, 718*). Rara in umbrosis Izamal, Nov. (930).

**BRADBURYA VIRGINIANA** (L. Bth) Kuntz Rev. Gen., i, 163.

Yucatan, *Johnson sine num.* specimen in Herb. Torrey Columbia Univ. "Vine, 20 feet, common in brush and forest lands about Izamal, its purple flowers appearing from August until December" (833). Yucatan, loc. ignot. (*Schott, 867*).

**CÆSALPINIA BONDUCELLA** (L.) Flem. As. Ris., xi, 159. *Cojones del Gato*. Celestun, near the sea, May 13, 1865 (*Schott, 402*).**CÆSALPINIA PULCHERRIMA** (L.) Sw. Obs., 166.

Merida, "fragrant," July 25, 1865 (*Schott, 570*). Merida, Feb. 2, 1864 (*Schott, 178*, in Herb. U. S. Nat'l Mus.). Silam (681).

**CALLIANDRA GRACILIS** Klotz. Saund. Refug. Bot., t. 294.

"An uncommon shrub 20 feet high, at Progreso" (1178). Merida, August, 1865 (*Schott, 844*).

**CALLIANDRA PORTORICENSIS** (Jacq.) Bth. Lond. Jour. Bot., iii, 99. **Pich.**

Near Hacienda Uislimchac, April 2, 1866, leaves and flowers undeveloped (*Schott, 769*).

**Canavalia ensiformis** (L.) de C. Prod ii, 404.  
Merida, Feb. 1, 1866, in fruit only (*Schott*, 729).

*Haba.*

**Cassia\* sp.**

"An interesting plant allied to *C. biflora* L, but indeterminable in the absence of fruits. The ash-colored twigs and very small leaflets do not correspond with any existing species of that group" (C. L. P.) Leaves alternate fascicled, common rachis 5-8 lines long, leaflets 3-jugal, 1-3 lines long, varying from mucronate or truncate to emarginate, stipules geminate, aristate; inflorescence solitary, sparse, from the terminal leaf fascicles; peduncle attenuate, longer than the leaf, jointed at about the middle, flowers large 4-7 lines long. Scrub lands near Sisal, May 6, 1865 (*Schott*, 414).

CASSIA BIFLORA Linn? Sp. Pl.; 378.

"Probably this species, a comparison with the type is, however, necessary" (C. L. P.) Tizimin, March 28, 1866 (*Schott*, 982).

CASSIA EMARGINATA Linn. Sp. Pl., 346.

**Xtuab.**

Without locality *Valdez* (7). North of Merida on the road to Progreso, April 7, 1865 (*Schott*, 261). Tunkas, tree 20 feet, flowers orange, March 3, 1890 (*Stone*, 236, in Herb. Acad. Sci., Phila.).

**Cassia flavicoma** H. B. K., Nov. Gen. et. Sp. vi. 366.

"A rare herb, 3 feet high, in the forests of Buena Vista Xbac, flowering in September" (1113).

**Cassia hirsuta** Linn. Sp. Pl., 540.

**Salché.**

"A common herb, 4 feet high, in waste places and on farms and open lands about Izamal, producing its chrome-yellow flowers from July to December" (757, 1090).

**Cassia leptocarpa hirsuta** Benth. Trans. Linn. Soc. Lond., xxvii. 531.

"A shrub 6 feet high, common near cenotes about Izamal, the bright yellow flowers produced in October" (999); also "frequent in waste places about Izamal, where the plant often blooms several years in succession" (1016).

**Cassia Liebmannii** Bth. Trans. Linn. Soc., 27, 549.

Along road between Merida and Progreso, May 3, 1865 (*Schott*, 299).

CASSIA OCCIDENTALIS Linn. Sp. Pl., 377.

Nohcacab, Oct. 19, 1865 (*Schott*, 975).

\* Species determined by Mr. C. L. Pollard.

**Cassia polyphylla** Jacq. Coll. iv. 104. Ic. Rar. t. 460.

"In mimosetis prope Sisal" (*Schott, 414*). "Shrubby 6 feet high, in brush lands about Izamal, the golden-yellow flowers appearing throughout the year" (391). Tekanto, shrub 3 feet high, Feb. 25, 1890 (*Stone, 197, 219*, in Herb. Acad. Sci., Phila.).

**Cassia procumbens** Linn. Sp. Pl., 380.

On the downs at Progreso, May 4, 1865 (*Schott, 290*).

**Cassia racemosa** Mill. Dict. ed. 8, n. 19.

**Habinpec.**

"An abundant tree 60 feet high in brush and forest lands about Izamal, producing its large globular clusters of orange-yellow flowers from February to June" (354). Merida, Aug. 30, 1865 (*Schott, 974*). Tekanto, tree 20 feet high, flowers bright orange, Feb. 25, 1890 (*Stone, 215*, in Herb. Acad. Sci., Phila.).

CASSIA SERICEA SWZ. Fl. Ind. Occ., 724.

Merida, June 11, 1865 (*Schott, 440*).

**Cassia tristicula** H. B. K., nov. Gen. et Sp., vi, 367.

Ruins of Uxmal, Sept. 1865 (*Schott, 976*).

**Cassia villosa** Mill. Gard. Dict., ed. viii, 4.

**Salché.**

Suburbs of Merida, Nov. 25, 1864 (*Schott, 1*).

**Clitoria Mexicana** Lk. Enum. ii., 235.

"Vine 10 feet, common at Tekax" (1136).

**Cracca bicolor** Bull. Herb. Boiss ii., 444 t. xi.

"A common herb 4 feet high, in the forests of Buena Vista Xbac, in bloom in April" (1058).

CRACCA GREENMANII Millsp. Field Col. Mus., Bot. i. 299 t. xiii.

"An ascending herb 12 to 18 inches high, abundant along roadsides about Izamal, where it is almost without exception more or less browsed by cattle. The flowers appearing throughout the year are pale yellow with a purplish standard" (301). "Abundant near Merida" (*Veldez, 69*). Tekanto, Feb. 25, 1890 (*Stone, 205, 209*, in Herb. Acad. Sci. Phila.).

CRACCA VILLOSA CINEREA (L.) Kuntze, Rev. Gen. 1. 174.

Merida, Aug. 3, 1865 (*Schott, 537*).

CROTALARIA PUMILA Ortega Hort. Matr. 23.

Flats about Nohpat and Nohcacab, Nov. 1865 (*Schott, 715*).

The legume in these specimens is shortly pediceled, not sessile as noted in description. Our specimens agree fairly well with Müller Herb. Mex., 664, 1745; and especially with Buslamente y Rocha's 19 in the Torrey Herb.

**Dalea diffusa** Moric. Pl. Nouv. Am. 8. t. 6.

"Abundant in old fields and on town lots near Izamal, Nov. Height 4 feet. Flowers a very evasive royal-purple, which, seen at a short distance, seems a glimmer of light between blue and crimson, always attractive" (934).

In general habit our plant agrees with Pringle (1793) Jalisco, 1888, and Palmer (507) Acapulco 1894-5, except that in the former the leaflets are smaller, narrower, more crowded and nearer entire; and in the latter still smaller than our form or the Jaliscoan. Leaves in our form 1.75 to 2.25 inches long, leaflets .4 inch long, .1 inch wide, with glandular punctæ strongly marked.

**DESMANTHUS DEPRESSUS** H. & B. Willd. Sp. iv. 1046. **Camba-pich.**  
 "Ten feet high, rare about Izamal, its white flowers appearing in October" (1018).

**Diphysa Carthaginensis** Jacq. Enum. Pl. Carib., 28.  
 About Merida, March 16, 1865 (Schott, 247), April 20, 1865 (Schott, 505, in Herb. U. S. Natl. Mus.).

**Diphysa robinoides** Benth. ex Bth. & Oerst. l. c. 11. **Quouc.**  
 "An abundant tree 60 feet high in the brush and forest lands near Izamal, where it opens its bright golden-yellow flowers in February while the leaves are immature" (344). Yucatan, without locality (Valdez, 58).

**Erythrina Corallodendron** Linn. Sp. Pl. 706.  
 Shrub 5 feet high, with sharp, mostly recurved prickles and crimson flowers, Tekanto, Feb. 25, 1890 (Stone, 217, in Herb. Acad. Sci. Phila.).

**ERYTHRINA CORALLOIDES** Mocq. & Sesse, ex de C. Prod. ii, 413.  
 Merida, Aug. 28, 1865 (Schott, 615). Loc. ignot. (Schott, 921).

**Galactia multiflora** Robn. Proc. Am. Acad. xxix, 315.  
 "Vine 20 feet, abundant in brush and forest lands about Izamal, producing its mottled flowers from September to March" (566). Also collected in conjunction with 498 in similar habitat.

**INDIGOFERA ANIL** Linn. Mant. 272. *Anil.* **Choh.**  
 "Shrubby, 4 to 8 feet high, abundant in all northeastern Yucatan, where it was at one time extensively cultivated, and whence Indigo was exported as late as 1885. To-day (1896) not one plantation exists in the neighborhood of Izamal, which place was once the center of its greatest production" (1057). Merida, Oct. 25, 1865 (Schott, 883).

**Meibomia albiflora** (Salzm) Kuntze, Rev. Gen. i, 197.  
 Yucatan, loc. ign. (Johnson, 36<sup>b</sup>, Visa in Herb. Torrey, Columbia University).

**Meibomia Neo-Mexicana** (A. Gr.) O.K., Rev. Gen. i, 198. **Kintah.**  
 Chuburná, Oct. 10, 1865 (Schott, 866); herb 2 feet high, common in the forests of Tekax, September (1130); herb 10 inches, forests of Buena Vista Xbac (1118). Schott's plant and Gaumer's 1130 are glabrous, while 1118 is finely pubescent.

**MEIBOMIA SCORPIURUS** (Sw.) Kuntze, Rev. Gen., 198.

"Vine 4 feet, common in shady places near Izamal, October to January" (929).

**Mimosa asperata** Linn. Syst. ed x. 1312.

Yucatan, loc. ign. (*Johnson, 40, Visa* in Herb. Torrey, Columbia University).

**Mimosa somnians** Humb. & Bonpl. ex Willd. Sp. Pl. iv, 1036.

Yucatan, loc. ign. (*Johnson, 35<sup>b</sup>, Visa* in Herb. Torrey, Columbia University).

**MUCUNA PRURIENS** (L.) de C., Prod. ii, 405.

Suburbs of Merida, Dec. 11, 1864, in fruit only (*Schott, 155*).  
"Vine 25 feet, common on brush lands and old fields, frequently also in forests about Izamal" (918).

**Myroxylon peruiferum** Linn. f. Suppl., 233.

**Naba.**

Yucatan loc. ignot. (*Schott, 804, in Herb. U. S. Natl. Mus.*).

**PAROSELA DOMINGENSIS** (de C.).

*Dalea Domingensis* de C. Tekanto, Feb. 27, 1890, 3 feet high, flowers dull red (*Stone, 193, in Herb. Acad. Sci. Phila.*).

**Phaseolus dysophyllus** Benth. Pl. Hartw. 287.

"Vine 20 feet, common in the brush and forest lands about Izamal" (923).

**PHASEOLUS LUNATUS** Linn. Sp. Pl. 1016.

"Vine 10 feet, growth frequent in the forest lands about Izamal" (1059).

**PISCIDIA ERYTHRINA** Linn. Syst. ed. x, 1155.

**Habi.**

Yucatan loc. ignot. (*Schott, 260*).

**Prosopis juliflora** (Swz.) de C. Prod. ii, 447.

**Xcaimek.**

"Tree 40 feet high, common at the Port of Silam, producing its greenish-white flowers in April" (641).

Foliage much denser and leaflets larger than in the north Mexican and North American forms.

**Pterocarpus Draco** Linn. Sp. Pl. 1662.

**Subinché.**

"Tree 80 feet high, common in forests about Izamal, developing its yellowish-brown flowers in February" (377).

**RHYNCHOSIA MINIMA** (L.) de C. Prod., ii, 385.

Ruins of Nohpat, Nov. 24, 1865 (*Schott, 718*).

**Sesbania grandiflora** Poir. Dict. vii., 127.

*Pico de Flamingo.*

"Shrub 15 feet high, naturalized at the Port of Silam, producing its large crimson flowers in April" (670). Hacienda Chucá cult? March, 1866 (*Schott, 763*).



**Sesbania macrocarpa** Muhl. ex. Raf. Fl. Lud. 137.

“Herb 4 feet high, infrequent on waste lands about Izamal, flowering in September” (907).

**Stylosanthes biflora** (L.) B. S. P.

“Frequent at the margin of a large sarteneja on the Cantoila road northwest of Izamal” (908).

TAMARINDUS INDICA L., Sp. Pl., 48.

*Tamarindo.*

Merida, May 20, 1865 (*Schott, 356*).

#### OXALIDACEÆ.

OXALIS BERLANDIERI Torr. Bot. Mex. Bd., 41.

Hacienda, Cucul, Sept. 8, 1865 (*Schott, 625*).

OXALIS LATIFOLIA H. B. K. Nov. Gen. et Sp., v., 237.

Merida, June 16, 1865 (*Schott, 490*). Aguada and Cenote Nabalá, Aug. 8, 1865 (*Schott, 918*).

#### MALPIGHIACEÆ.

**Bunchosa glandulosa** (Cav.) de C. Prod., i., 581.

**Sipché.**

“Shrub 20 feet high, common in the forests about Izamal, the bright yellow flowers produced from February to June” (411). Suburbs of Merida, April 14, 1865 (*Schott, 84*). Tekanto, 6 feet high, Feb. 27, 1890 (*Stone, 192*, in Herb. Acad. Sci. Phila.).

**Gaudichaudia flipendula** Juss. *infra* iii., 594.

**Chilillo-ak.**

“Common at Progreso, October” (1138).

HETEROPTERYS BEECHEYANA A. Juss. Arc. Mus. Par., iii., 475.

In fruit only, Merida suburbs, Dec. 11, 1864 (*Schott, 158*).

**Heteropteris Yucatanensis** sp. nov.

Frutex, foliis oppositis ovato-lanceolatis acutis e basi cuneatis integris, petiolis brevis eglandulosis, ramis ad apice tomentosis, paniculis terminalibus paucifloris, pedicellis ultra medium articulatis, bracteis deltoideis minimis, calycis 8-glandulis albo-tomentosis. Ramis brevis 4-6 poll. long, foliis 1-2.5 poll. long, .5-1 poll. lata; petiolis 1-3 lin. long. Frutex 15 pedalis, in terræ inutiliæ. Izamal, Sept. (816).

**Stigmaphyllon lupulus** Wats. Proc. Am. Acad., xxi., 461.

“Vine 15 feet, infrequent on old stone fences about Izamal, producing its bright lemon-yellow flowers from February to July” (408). Our specimens agree best with Pringle’s 4102 from Tamasopo, San Louis Potosi, Mexico, 1892; the leaves varying from orbicular-cordate through ovate-cordate and simply hastate to deeply hastate-lobed. At Citas, March 31, 1866 (*Schott, 771*). Tunkas, March 3, 1890, a woody climber (*Stone, 230*, in Herb. Acad. Sci. Phila.).

**Tetrapteris inæqualis?** Cav. Diss. 9, t., 260.

“Uncommon in forests and brush lands near Izamal” (972).

### ZYGOPHYLLACEÆ.

TRIBULUS MAXIMUS Linn. Sp. Pl., 387.

Merida, Feb'y 8, 1865; and March 3, 1865 (*Schott, 25, 97*).

TRIBULUS TERRESTRIS CISTOIDES (L.) Oliver.

Downs at Progreso, April 6, 1865 (*Schott, 285*).

### RUTACEÆ.

ZANTHOXYLUM PTEROTA H. B.K. Nov. Gen. et Sp., vi., 3. **Xic-ché.**

Merida, June 11, 1865 (*Schott, 450*).

### SIMARUBACEÆ.

ALVARADOA AMORPHOIDES Liebm. Kjoeb. Vid. Med., 1853, 100.

Suburbs of Merida, Dec., 1864 (*Schott, 151*). Tekanto, tree 20 feet high (*Stone, sine num.*, in Herb. Acad. Sci. Phila.).

**Simaruba glauca** de C. Ann. Mus. Par., xvii., 323. **Xpaxakil.**

“Tree 100 feet, common in the forests about Izamal, producing its light green flowers in February” (439).

SURIANA MARITIMA Linn. Sp. Pl., 284.

**Xpanoil.**

On the downs at Progreso, April 4, 1865 (*Schott, 306*).

### BURSERACEÆ.

**Protium heptaphyllum** (Aubl.) Mch., Kjoeb. Vid. Med., 1873, 55.

*Copal.* **Pom.**

Near the hacienda Cucá, March 26, 1866 (*Schott, 757*).

### MELIACEÆ.

CEDRELA ODORATA Linn. Syst. ed., x., 940.

Merida (*Schott, 199*, in Herb. U. S. Natl. Mus.).

**Trichilia spondioides** Jacq. Pl. Carib., 20.

**Xkulinsis.**

“Tree 40 feet high, common in the forests about Izamal, producing its pale green flowers in June” (712).

### EUPHORBIACEÆ.

ACALYPHA ALOPECUROIDEA Jacq. Ic. Rar., iii., 19.

“Herb 16 inches high, common in shaded places near Izamal, flowering from July to Dec.” (1017). Chocholá, Sept. 13, 1865 (*Schott, 44*).

ACALYPHA SETOSA A. Rich. Fl. Cub. Fan., ii., 204.  
Merida, 1865 (*Schott, 480, 968*).

**Acalypha Yucatanense** sp. nov.

**Miscax.**

*Rhombifoliae.* Spicis ovoideo-cylindricis breviuscule pedunculatis, bracteis fœm. densis, 1-floris, late obovatis, hirsutis, 7-9 dentatis, dentibus triangularis acutis, calycis fœm. laciniis ovatis acutis hirsutis, ovario tuberculato et hispido, stylis 3 tenuibus, seminibus lævibus ecarunculatis. Herba pilosa 20 pollicaria, caule erectis simplicis vel inferne dubiter ramosis, limbus foliorum .5-1 poll. long, ovatis acutis crenulato-serratis e basi deltoideis, supra et infra pilosus, petiolum limbum longiorum. Ad Progreso (1176).

**Codiæum variegatum** Blume Bijd. 606.

Merida, at Quinta del Obispo, cult? March 6, 1865 (*Schott, 244*). Merida, cult. April 1887 (*Millsbaugh, 24*).

**Croton ciliato-glanduliferus** Ortega, Hort. Matr. 51.

Tekanto, Feb. 27, 1890 (*Stone, 216*, in Herb. Acad. Sci., Phila.).

CROTON CORTESIANUS H. B. K. Nov. Gen. et. Sp., ii., 83.

Merida, Nov. 1864 (*Schott, 172*). Tekanto, shrub 5 ft., Feb. 25, 1890 (*Stone, 201*, in Herb. Acad. Sci., Phila.).

CROTON FLAVENS Linn. Pl. Jam. Pug. 28.

Cenoté de Dambula near Merida, Jan. 20, 1865 (*Schott, 172*).

CROTON LOBATUS Linn. Sp. Pl. 1427.

Ruins of Uxmal, Sept. 15, 1865 (*Schott, 955*).

CROTON HUMILIS Linn. Syst. ed x., 1276.

Cenote de Dambula, Merida, Jan. 20, 1865 (*Schott, 172*).

CROTON MARITIMUS Walt. Flor. Carol., 239.

Progreso, April 4, 1865 (*Schott, 209*, in Herb. U. S. Natl Mus.).

DALÉCHAMPIA SCANDENS Linn. Sp. Pl., 1423.

Merida, June 30, 1865 (*Schott, 514*).

**Xmoolcoh.**

EUPHORBIA ADENOPTERA Bertol. Misc. Bot., iii., 436.

"Prostrate herb spreading one foot. Chaltumha road, October" (938). The densely floral and folial form with all characters closely aggregate.

"Semi-erect 10 inches high, Sitalpech road, November" (939). The more open ascending form. Sisal, Nov. 1865 (*Schott, 966*).

**Euphorbia arenaria** (Kth.) H. B. K. Nov. Gen., ii., 46.

Merida at Quinta del Obispo, Nov. 23, 1864 (*Schott, 15*, in part).

**EUPHORBIA ARMOURII** Millsp. Field Col. Mus., Bot., i., 28.

“Herb 18 inches high, uncommon, found only at the mouths of caves, and in very damp, shady places near Izamal, November” (975). Of larger growth, and with young parts more densely hirtellate than in the type. Merida, Cenoté Chambula, Dec. 14, 1864; Quinta del Obispo, Nov. 22, 1864 (*Schott, 76*).

**EUPHORBIA BUXIFOLIA** Lam. Encyc., ii., 421.

Seaside at Progreso, April 4, 1865 (*Schott, 308*).

**Euphorbia dentata** Michx. Fl. Bor. Am., ii., 211.

Merida, at Quinta del Obispo, Nov. 23, 1864 (*Schott, 15*).

**Euphorbia Gaumerii** sp. nov.

**Sacio.**

Caule glabræ, lignosæ, scandenti, ternatim ramoso, ad articulationes incrassata, ramis longe articulatis plus minusve pilis strigosis sparsis, foliis oppositis ovato-lanceolatis obtusis vel acutis e basi cuneata, petioli quaterne longiori, glandulis stipularibus brevis, cymis ad articulationes ramularum corymboso-paniculatis, polycephalis, ad apice racemoso-paniculatis, tota laxiuscule, pedunculatis divaricatim dichotomis; foliis floralibus rarius minimis ovato-lanceolatis, involucris pedicillatis turbinatis adpresse lanatis, lobis spathulatis irreguliter dentato-fimbriatis, glandulis transverse sub-plicatis appendice triplo latiori orbiculata minute serrato-dentata, stylis brevibus, stigmatibus crassis profunde bi-partitis. Capsula (juniori) glabra, profunde sulcata, coccis obtuse carinatis, semina. . . . . Frutex scandentis 10 pedalis. Folia 1-2 poll. longa, 9-12 lin. lata, cymæ laxiuscule oppositæ. Involucris lin. longa, duo-lineam lata. Species, prox. *colletioidea* Bth. (alectoroctonum). In sylvis ad Buena Vista Xbac (1073).

**Euphorbia graminea virgata** var. nov.

**Onob kax.**

“Herb 2 feet high, common in moist places near Tekax, September” (1128).

Differs from the species in having longer and virgate branches, larger leaves, smaller involucral lobes, broader and blunt pointed appendages having a tendency to trilobation, and smaller seeds with fewer scrobiculæ arranged in two transverse lines, sharper rugæ, a sharper triangular apex, and no median line.

**Euphorbia graminea lancifolia** var. nov.

“Herb 18 inches high, rare, railroad bank, northwest of Izamal, November” (969).

Branches less quadrangular and not hairy on the angles, lower ovate leaves repand dentate, upper leaves lanceolate, the floral linear on attenuate branchlets. Glandular appendages ovate, seeds longer and more sharply quadrangular than in *var. virgata*, and with three lines of pits.

**EUPHORBIA HETEROPHYLLA** Linn. Am. Acad., iii., 112.

“Herb 3 feet high, along old fence rows and in old fields about Izamal, in flower from September to December” (903). The usual form of the species with cuneate-lanceolate irregularly serrate leaves, smooth capsules, and virgate sparsely leafy branchlets.

Merida, Sept. 30, 1865 (*Schott, 960*). “This plant is used for washing clothing.”

**Euphorbia heterophylla** *forma*.

“Herb 4 feet high, common at Buena Vista Xbac, September” (1109).

A peculiar form in which the stem leaves are broadly lanceolate, entire or minutely and distantly dentate-serrate, strigosely short-hairy beneath; while those of the extended virgate branchlets are elongated-linear, quite resembling those of *var. graminifolia* Engelm. This form, but with some of the lower leaves, panduriform (*E. heterophylla cyathophora* (Jacq.) Boiss), was collected in 1879 at Fountaindale, Oregon, and Byron, Illinois, U. S. A., by the late Mr. M. S. Bebb.

**EUPHORBIA HETEROPHYLLA CYATHOPHORA** (K. and G.) Boiss de C. Prod., xv., 262.

Merida, 1865 (*Schott, 961*). This form of a very varied subspecies is completely vested with cyathiform leaves, from those of the base to the floral at the apex. Generally the basal leaves only distinguish the variety, the apical being either lanceolate or linear. The nearest counterpart of this plant was grown by Dr. Geo. Engelmann from seed sent him from Texas. (See sp. in U. S. Nat. Herb.)

**Euphorbia lasiocarpa** Klot. Nov. Act. xix, Supl. 414.

Merida, Aug. 23, 1865 (*Schott, sine num.*).

**EUPHORBIA MAYANA** Millsp. Field Col. Mus. Bot., I., 304.

“Open grounds at Buena Vista Xbac, April” (1039). Differs from the type collected near Izamal, only in having somewhat deeper and more irregular dentations in the margin of the appendages. Tekanto, shrub 5-8 feet high, branches short, horizontal, flowers yellowish-white, Feb. 25, 1890 (*Stone, 211*, in Herb. Acad. Sci. Phila.).

**Euphorbia pilosula** Engelm in litt.

“Herb 8 inches, common in the forests of Buena Vista Xbac, August” (1103).

**EUPHORBIA PILULIFERA PROCUMBENS** (de C.) Boiss. de C. Prod, xv., 21.

Merida, at Quinta del Obispo, Nov. 24, 1864 (*Schott, 56*). Merida, 1865 (*Schott, 967*). “Prostrate herb, abundant in shady places near Izamal, November” (1003).

**Euphorbia pulcherrima** (Grah.) Willd. Allg. Gart. ii., 27.  
Merida, at Quinta del Obispo, Jan'y, 1866 (*Schott*, 727).

EUPHORBIA RHYTISPERMA Engelm in Kl. et Gke., p. 34.  
"Waste grounds at Tekax, September" (1123). Larger and fuller growth than specimens collected at Izamal in 1895.

EUPHORBIA SERPYLLIFOLIA Pers. Ench. bot., II, 14.  
Intermixed with *E. pilulifera procumbens* collected by *Schott*, 967, at Merida, 1865.

**Euphorbia Xbacensis** sp. nov.

Glabra, caule erectis, quadrangulatis dichotomis, foliis petiolatis integris ovato-lanceolatis acutis e basi cuneata, summis minute hirtis diminutis lanceolatis, involucris minimis extus adpresse hirtis, lobis quadratis fimbriatis, glandulis 4, transverse avatis, appendice angustissime orbiculatis, margine integris vel obsolete lobulatis, stylis profunde bipartitis apice crassis; capsulæ hirtis profunde sulcatis, coccæ obtuse carinatis, semina grisea, ovate, reguliter foveolate albo in profundis. Bipedalis. Folia inferiora nulla, media semipollicaria longa, bilinea lata, supra minima.

§ *Cyttarospermum* prox. *E. Acerense*. Rara ad Buena Vista Xbac (1108).

**Gymnanthes lucida** Sw. Prod. Veg. Ind. Occ., 96.  
Merida, April 2, 1865 (*Schott*, 283, 316, in Herb. U. S. Natl. Mus.).

**Hura polyandra** Baill. Etud. gen. Euph., 543. *Jabilla*.  
A small tree 40 feet high, common at Progreso (1144).

JATROPHA CURCAS Linn. Sp. Pl., 1429. **Xcocal ché**.  
Merida, May 28, 1865 (*Schott*, 395).

JATROPHA URENS Linn. Sp. Pl., 1429. **Chaya**.  
Cercanias, Merida, April 30, 1865, and November (*Schott*, 496, 700).

JATROPHA URENS STIMULOSA (Michx.) Müll. de C. Prod., xv., 1101. **Xoat**.  
Merida, Nov. 5, 1865 (*Schott*, 700). Merida, in mimosetis et cercanias, April 12, 1865 (*Schott*, 496). "Herb 6 feet high. Common everywhere. Buena Vista Xbac, April" (1069). "The sting causes pain instead of burning or itching." Tunkas common, 4 feet high, March 3, 1890 (*Stone*, 232, in Herb. Acad. and Sci. Phila.).

**Manihot Carthaginensis** (Jacq.) Muell. D. C. Prod., xv., 1073. **Xhac ché**.  
"Herb 8 feet high, common along roads, and on waste lands, near Izamal" (1142). Leaflets nearly panduriform, strongly apiculate. Merida, June 25, 1865 (*Schott*, 518). The same panduriform lobed form.

**Manihot Manihot** (L.) Cock. Bull. Torr. xix., 1892. *Jucca*.  
Cultivated near the ruins of Kabah, Nov. 26, 1865 (*Schott, 957*).  
A form with 7 narrowly lanceolate, almost linear, leaf lobes, not  
in flower or fruit.

**Manihot rhomboidea** Muell. Arg. in Linn., 34, 205. **Chaché**.  
Merida, July 11, 1865 (*Schott, 518*). Dr. Schott spells the Maya  
name Xcaxe.

**PEDILANTHUS ITZÆUS** Millsp. F. Col. Mus. Bot., 1, 305. **Yaxhalal ché**.  
Merida, April 10, 1866, in full leaf (*Schott, 552*).  
(This species also appears in U. S. Natl. Herb., from Santa  
Domingo, collected by Wright, Parry and Brummel, without  
number.)

**RICINUS COMMUNIS** Linn. Sp. Pl., 1430.  
Merida, Feb'y 13, 1865 (*Schott, 241*). Merida, April 14, 1887.  
(*Millsbaugh, 46*).

**TRAGIA NEPETÆFOLIA** Cav. Ic., vi., 37. **Popox**.  
Merida, Aug. 20, 1865 (*Schott, 588*); February 11, 1865 (*Schott,*  
*448*). "The Mayas rub their limbs with this plant when suffer-  
ing pain."

## ANACARDIACEÆ.

**Anacardium occidentale** Linn. Sp. Pl., i., 283. *Maranon*.  
Hacienda Tahchebilá, Cult? March 31, 1866 (*Schott, 765*).

**Astronium** sp.  
"A rare tree 100 feet high." In fruit without leaves, Xcholac,  
March (582).

**MANGIFERA INDICA** Linn. Sp. Pl., 200. *Mango*.  
Merida. Cult? March 14, 1866 (*Schott, 815*). In these speci-  
mens the panicle is not pubescent, indeed only very sparsely  
hairy.

**Spondias lutea** Linn. Sp. Pl., Ed. ii., 613. *Cirueta*. **Mompin**.  
Suburbs of Merida, cult? April 4, 1865 (*Schott, 797*).

**Spondias purpurea** Linn. loc. cit. *Cirueta*. **Chiabal**.  
"Tree 30 feet high, very abundant in cultivation, the bright-  
pink flowers appearing in March" (432). "*Juntura*, flowers  
red" (704).

## CELASTRACEÆ.

**Maytenus phyllanthoides** Benth. Bot. Voy. Sulph., 54.  
"Shrub 10 feet, abundant at the port of Silam, April" (643).

## HIPPOCRATEACEÆ.

**Hippocratea obcordata** Lam. Illust., i., 100.  
Downs at Progreso, April 6, 1865 (*Schott, 317*).

## SAPINDACEÆ.

- Bumelia buxifolia** Willd. H. B. K. Nov. Gen. et Sp. t., 147.  
Downs at Progreso, April 7, and Celestun, May 12, 1865  
(Schott, 313).
- Bumelia glomerata** Griseb. Mem. Am. Ac., 1863, 518. **Puomucuy.**  
Suburbs of Merida, April 14, 1865 (Schott, 341). "Shrub 15  
feet high, abundant in brushlands about Izamal, producing its  
cephalic whorls of greenish-yellow flowers from March to  
April" (473).
- Bumelia microphylla** Griseb Plant. Cat. Cub., 165.  
Merida, March, 1865 (Schott, 341). "Shrubby, 10 feet high,  
common on brushlands near Izamal, April" (559, 469 in part).
- Paullina fuscescens glabrescens** Radlk. Mon. Sap., 56. **Kexak.**  
About Merida, Dec. 10, 1864 (Schott, 10, in Herb. U. S. Natl.  
Mus.).
- Serjania polyphylla** Poir. ex. Steud. Nom., ed. 2, ii., 568.  
About Merida, Dec. 10, 1864 (Schott, 10, part of the sheet  
upon which the previous species is mounted in Herb. U. S.  
Natl. Mus.).

## RHAMNACEÆ.

- Colubrina ferruginosa** Brong. Ann. Soc. Nat. x. 369. **Yax-Pukim.**  
A small tree 30 feet high, Mascab Pixoy, Jan. 22, 1895 (*Armour  
Exped.*, 157, 181). "Small tree 30 feet high, abundant in scrub  
and brushlands about Izamal" (355). Suburbs of Merida,  
"Oulubmay," April to November (Schott, 13, 331).
- Gouania Domingensis** Linn. Sp. Pl. 1663. **Xomak.**  
"Vine 30 feet, infrequent in the forests about Izamal, Novem-  
ber" (959). Merida, Nov., 1865 (Schott num. amiss. 658, in Herb.  
U. S. Natl. Mus.).

- Karwinskia Humboldtiana?** Zucc Nov. Stirp., i., 351.  
"Tree! 30 feet high, common from Merida to Progreso" (1171).  
Although our plant seems to properly fall under this genus and  
species; yet the ovary is 2-celled with each cell 2-ovuled, and the  
leaves, though opposite, are not pellucid-dotted.

## TILIACEÆ.

- CORCHORUS SILIQUOSUS** Linn. Sp. Pl. 746.  
"Shrub 3 feet high, common at Progreso" (1168). Aguada de  
Uxmal, Sept. 17, 1865 (Schott, 104, in Herb. U. S. Nat. Mus.).  
Differs from the description in the following characters: More  
shrubby than suffrutescent; branchlets with a double hairy line;  
flowers single, axillary, scattered along the branches.
- Lueha speciosa** Willd. Geo. Nat. fr. Neu. Schr. iii., 410. **Chacah.**  
At the Hacienda Chacah near Ticul, Oct. 21, 1865 (Schott, 679).



- Muntingia Calabura** Linn. Sp. Pl. 728. *Capolin.*  
 "Hacienda San Rafael de Xteppen southeast of Uman, Aug., 1865. Hacienda Saragossa, Sept. 9, 1865" (*Schott, 554*). "Tree 30 feet high, common at Progreso and Merida" (1163).

## MALVACEÆ.

- Abutilon crispum** Sweet, Hort. Brit. ed. i., 53.  
 "Merida at Quinta del Obispo, Nov. 20, 1864" (*Schott, 83*, in Herb. U. S. Natl. Mus.); suburbs of Merida, Dec. 3, 1864" (*Schott, 167*). "Herb 5 feet high, common on open lands near Izamal, flowering from August to December" (872). Calcehtoh, March 11, 1890, flowers yellow (*Stone, 271*, in Herb. Acad. Sci., Phila.).
- Abutilon incanum** (Lk.) Sweet, Hort. Brit. ed. i., 53.  
 "Herb 4 feet high, abundant in old fields and waste places near Izamal" (954). Our form agrees best with Palmer's 1281 from Armeria, Mex., 1891.

- Abutilon sidoides** Hemsl. Diag. Pl. Nov. 24. **Xacmixbil.**  
 "Shrubby, 4 feet high, common at Tekax, September" (1133). Our specimens differ from the type in having somewhat longer petioles, and in being more branchy like Pringle's 4583 Jalisco, 1893.

- ANODA HASTATA** Cav. Diss. i., 39.  
 Merida, Nov. 28, 1864 (*Schott, 12*). Hacienda Saragossa, Sept. 8, 1865 (*Schott, 353*, in Herb. U. S. Natl. Mus.).

- Anoda parviflora** Cav. Ic. v., 19. **Jaiunoi.**  
 "Herb 3 feet high, abundant in old fields near Izamal, producing its pale-yellow flowers from September to October" (821).

- Anoda triangularis** de C. Prod. i., 459.  
 Fence rows in the suburbs of Uman, April, 1887 (*Mills-paugh, 19*).

- Gossypium Barbadense** Linn. Sp. Pl. 975. **Xchup.**  
 Suburbs of Merida, Aug. 24, 1865 (*Schott, 602*) Calcehtoh, March 11, 1890 (*Stone, 269*, in Herb. Acad. Sci. Phila.).

- Gossypium herbaceum** Linn. Sp. Pl. 975. *Algodon.*  
 "Herb 10 feet high, often becoming a 15 foot shrub. Extensively cultivated in Yucatan during the civil war in the United States, now only on a small scale for domestic use" (935). Our specimens are of the form *G. hirsutum* L. and agree with Berlandier's Texano-Mexican 313, and Simpson's Florida specimens of 1889.

- Gossypium religiosum** Linn. Syst. ed. xii. 642.  
 Suburbs of Merida, August, 1865 (*Schott, 602*).

**Hibiscus elatus** Sw. Fl. Ind. Occ. ii, 1218. **Xholó.**  
Hacienda Guayalú, cult? Sept. 19, 1865 (*Schott, 629*).

**Hibiscus mutabilis** Linn. Sp. Pl. 977. *Flor de Cortejo.*  
Cultivated at Merida, Feb. 16, 1866 (*Schott, 744*).

**HIBISCUS TUBIFLORUS** Moc. & Sesse, de C. Prod. i. 447.  
Tunkas, 3 to 4 feet high, flowers rose-red, March 3, 1890 (*Stone, 257*), in Herb. Acad. Sci. Phila.).

**Malachra capitata** Linn. Syst. ed. xii. 458.  
At the Aguada de Uxmal, Sept. 17, 1865 (*Schott, 883*).

**Malvastrum tricuspdatum** A. Gray, Pl. Wright i. 16. *Malva.*  
*M. Coromandelianum* Garcke. "Shrubby, 4 feet high, abundant in old fields near Izamal, October" (1020). Our plant agrees with that of *P. Sintenis* from Porto Rico.

**MALVAVICUS ARBOREUS** Cav. Diss. t. 48.  
Two quite distinct forms of this species are found in our region: The one with small obtuse slightly and irregularly crenate leaves ( $\frac{1}{2}$  to  $1\frac{1}{4}$  inch long by  $\frac{3}{8}$  to  $\frac{1}{2}$  inch broad), collected only along the coast. (Gaumer, Silam, April, 1895, *Schott* Downs at Progreso, April, 1865, 271); the other with large acute crenate leaves (1 to 3 inches long by  $\frac{3}{4}$  to 2 inches broad), found inland (Gaumer 580, *Millsbaugh, Armour Exped. 60, Schott*. Merida, Jan. 1865, 177; and Ruins of Uxmal, Sept. 15, 1865, 643). The leaves of the form from the coast have a few scattered pilose hairs beneath, those of the inland form scattered stellate hairs, while in the specimens from the more humid and higher region of Uxmal the leaves are densely woolly.

**SIDA ACUTA CARPINIFOLIA** (L.f) K. Schum Fl. Braz. Fasc. cix. 326.  
"Herb 4 feet high, frequent in brushlands about Izamal (902); and in the forests about Buena Vista Xbac, April" (1054). Tekanto common, Feb. 27, 1890 (*Stone, 194*, in Herb. Acad. Sci. Phila.).

**Sida anomala** St. Hil. Fl. Bras. Mer., i. 177.  
Merida, Sept. 25, 1865 (*Schott, 116*, in Herb. U. S. Natl. Mus.).

**SIDA SUPINA PILOSA** (Cav.).  
*S. pilosa* Cav. Diss. i. 9. t. 8. Tekanto, Feb. 27, 1890 (*Stone, 196*, in Herb. Acad. Sci. Phila.).

**Wissadula mucronulata** A. Gray, Bot Mex. Bound 39. **Ounikax.**  
"Herb 5 feet high, common in old fields about Izamal, October" (968). Our plants agree with Pringle 4603 Jalisco, Mex., which have, however, a much lighter upper surface to the leaves.

**Wissadula tricarpellata** Robn. & Greenm.  
"Herb 2 feet high, common in open lands at Buena Vista Xbac, April" (1056).

## BOMBACACEÆ.

CEIBA CASEARIA Medic. Malv., 16.

Merida, Feb. 6, 1865 (*Schott, 204*). Tekanto, March 9, 1890 (*Stone, 224*, in Herb. Acad. Sci. Phila.).

CEIBA SCHOTTII Brit. & Baker. Jour. Bot. xxxiv, 173.

Merida, Aug. 1865 (*Schott, 205*), Type collection.

PACHIRA FASTUOSA (Sesse) Decne. Fl. des Serres, xxiii, 48.

Merida, Sept. 1865 (*Schott, 205*).

QUARARIBEA FIELDII Millsp. Field Col. Mus. Bot. i. 309. **Maha.**

Fine fruiting specimens from Merida at Quinta del Obispo Nov. 18, 1864 (*Schott, 18*). "The flowers are used to flavor chocolate."

Fruit ovoid rusty-downy, its lower two-thirds enveloped by the persistent calyx, the apex truncate-apiculate,  $1\frac{1}{4}$  inches long,  $\frac{3}{4}$  inch diameter.

## STERCULIACEÆ.

AYENIA PUSILLA Linn. Syst. ed. x, 1247.

"Herb 2 feet high. Abundant in open lands about Izamal, April" (1061).

**Ayenia** sp. nov? (**Yucatanense**).

"Herb 4 feet high. Common in forests at Buena Vista Xbac. In fruit but without flowers, April" (1052).

Foliis longe petiolatis ovato-lanceolatis, acuminatis e basi truncatis, grosse et irreguliter dentatis, strigose hirsutis capsula dense hirsutis, longi pedunculatis, semina orbiculatis rufo-maculatis. Folia  $1\frac{1}{2}$  -  $2\frac{1}{2}$  poll. long. 5-10 linea lata; petiola 1- $1\frac{1}{2}$  poll. long.

**Buettneria Carthaginensis** Jacq. Stirp. Am. Piot. 41. **Xtexak.**

"Shrub 15 feet high, common in forests about Izamal, putting forth its light-green flowers in August" (786). Our plants are much nearer the Bolivian form than that of other Mexican stations.

GUAZUMA POLYBOTRYA Cav. Ic. iii. 51.

**Pixoy.**

Suburbs of Merida, August to December, 1865 (*Schott, 148*). Merida, April 14 (*Schott, 342*, in Herb. U. S. Natl. Mus.).

HELICTERES BARUENSIS Jacq. Enum. Pl. Carib., 30.

**Outup.**

Merida, June 14, 1865 (*Schott, 397*). (792 in Herb. U. S. Natl. Mus.)

MELOCHIA TOMENTOSA Linn. Syst. Ed. x, 1140.

Merida, December, 1864 (*Schott, 608*). Ticul, flowers blue, March 12, 1890 (*Stone, 276*, in Herb. Acad. Sci., Phila.).

STERCULIA CARTHAGINENSIS Cav. Diss. vi, 353.

Borders of brushlands at Uman. April, 1887 (*Millsbaugh, 8*).

**Triumfetta semitriloba** Jacq. Enum. Pl. Carib. 22. **Ochmul.**  
Tizimin, March 28, 1866 (*Schott, 759*). "Shrub 10 feet high,  
infrequent in the brushlands about Izamal, November" (1005).

WALTHERIA AMERICANA Linn. Sp. Pl. 941.

*W. Indica* L. in prev. contrib. Charcos de Celestun, May 13,  
1865 (*Schott, sine num.*). Calcehtoh, March 11, 1890 (*Stone, 268*,  
in Herb. Acad. Sci., Phila.).

#### GUTTIFERÆ.

**Clusia flava** Jacq. Enum. Pl. Carib., 34. **Chunup.**  
Hacienda de Chablé, August 8, 1865, used as a remedy for  
syphilis (*Schott, 539, 924*).

#### BIXACEÆ.

**Amoreuxia palmatifida** Moc & Sesse de C. Prod ii, 638.  
"Shrub 4 feet high, frequent at Progreso" (1155).

MAXIMILIANA HIBISCOIDES (H. B. K.) Kuntze Rev. Gen. 44.

Between Citas and Tizimin, March 23, 1866 (*Schott, 752*). Tek-  
anto, March 9, 1890 (*Stone, 223*, in Herb. Acad. Sci., Phila.).

#### FLACOURTIACEÆ.

**Samyda serrulata** Linn. Sp. Pl. 558. **Puus Mucuy.**  
Near Merida, April 22, May 1, June 25 and 29, 1865 (*Schott,*  
*603*). "Tree 25 to 40 feet high, common in brushlands near  
Izamal, developing its white flowers from May to October" (699,  
858, 1063).

#### TURNERACEÆ.

TURNERA DIFFUSA APHRODISIACA (Willd) Urban, Jahrb. Bot. Gard.  
Berol ii, 127.  
Hacienda Saragossa, Sept. 9, 1865 (*Schott, 889*).

#### PASSIFLORACEÆ.

PASSIFLORA FOETIDA Linn. Sp. Pl. 959.

*P. Ciliata* Dryand. Ait. Hort. Kew. ed. 1, iii. 310. Calcehtoh,  
March 11; Ticul, growing over the church, March 16, 1890.  
Flowers white, filaments purple, pistils crimson, stigma green  
(*Stone, 270, 286*, in Herb. Acad. Sci., Phila.).

**Passiflora pulchella** H. B. K., Nov. Gen. et Sp. ii, 134 ?

"Vine 10 feet, uncommon in forests near Izamal, blossoming  
in September" (796).

**Passiflora suberosa** Linn. Sp. Pl., 958.

Merida, October 10, 1865 (*Schott, 898*).

## CARICACEÆ.

CARICA PAPAYA Linn. Sp. Pl., 1036.

Cult? Quinta del Obispo, Merida, March, 1865 (*Schott*, 245).

## LOASACEÆ.

GRONOVIA SCANDENS Linn. Sp. Pl., 202.

Merida, September 1, 1865 (*Schott*, 620).

## LYTHRACEÆ.

PUNICA GRANATUM Linn. Sp. Pl., 676.

Cult? Merida, May 22, 1865 (*Schott*, 40).

## RHIZOPHORACEÆ.

RHIZOPHORA MANGLE Linn. Sp. Pl., 634.

Yucatan ad Manglares, Nov. 9, 1864 (*Schott*, num. amiss.).

## MYRTACEÆ.

PIMENTA OFFICINALIS Lindl. Coll. Bot. t. 19.

*Pimiento.*

*Myrtus Pimenta* L., *P. vulgaris* Wight. Merida, cult? April 30, 1865 (*Schott*, 358).

PSIDIUM GUAJAVA Linn. Sp. Pl. ed. i, 470.

*Guayaba.*

Cult. Merida, March 23, 1865 (*Schott*, 257).

## COMBRETACEÆ.

BUCERAS CATAPPA (Linn.) Hitch. Mo. Bot. Gard. 1893, 85. *Almendra.*

*Terminalia* Linn. Hacienda San Rafael Xteppen, August 7, 1865; Hacienda Cucul, near Merida, September 8, 1865 (*Schott*, 550, 921); Ticul, March 13, 1890 (*Stone*, 280, in Herb. Acad. Sci., Phila.).

**Conocarpus sericea** Forst., Don. Gen. Syst., ii, 662. **Xtabché.**  
*Botoncillo.*

"A densely silky sericeous shrub 20 feet high, abundant on the borders of all salt marshes along the coast. Progreso" (1164) (*Schott*, 266, 267). This form is distinct from *C. erecta* in all its characters and should (according to our specimens) be kept apart.

**Gyrocarpus Americanus** Jacq. Select. Am. 282. **Xkis.**

"Tree 80 feet high, common in the forests about Izamal, producing its large cymose greenish flowers from March to June" (491, 530). Yucatan loc. ignot. (*Schott*, 530, in Herb. U. S. Nat'l Mus.).

## UMBELLIFERÆ.

**Hydrocotyle prolifera** Kellogg, in Proc. Calif. Acad. i, 15.

"Herb 4 inches high, abundant about the aguada of Xcholac, February" (445). Our specimen agrees with M. Bourgeau's 400 from the banks of a canal near Mexico City.

## LENNOACEÆ.

- Lennoa cœrulea** Fourn. Bull. Soc. Bot. Fr., xvi, 11.  
 "In the forests of Buena Vista Xbac, September" (1116).

## MYRSINACEÆ.

- JACQUINIA ARMILLARIS Jacq. Enum. Pl. Carib., 15. **Muyché.**  
 Seaside at Sisal, May 15, 1865 (*Schott, 357*). The same unarmed form described on p. 312, Contrib. ii; there is, however, an unnumbered specimen in Herb. U. S. Nat'l Mus., collected at Progreso, April 4, 1865, by Dr. Schott, in which the leaves are characteristically armed at the tip.

## PLUMBAGINACEÆ.

- Plumbago Capensis** Thunb. Prod. Pl. Cap. 33. *Embelezo.*  
 "Shrub 4 feet high, in general cultivation in towns on the peninsula" (1151).
- PLUMBAGO SCANDENS Linn. Sp. Pl., 215. **Xcabaac.**  
 Merida, June, 1865 (*Schott, 612*); Merida, Nov. 14, 1864 (*Schott, 29*, in Herb. U. Nat'l Mus.).

## SAPOTACEÆ.

- ACHRAS SAPOTA Linn. Sp. Pl., 470. **Ya.**  
 Merida, cult? March 23, 1865 (*Schott, 258*).
- CHRYSOPHYLLUM CAINITO Linn. Sp. Pl., 192. *Cainito.*  
 "Tree 30 feet high, common in the forests about Izamal, producing its cream-colored flowers in August" (896). Cultivated at Merida, Quinto del Obispo, Dec. 24, 1864 (*Schott, 72*).
- Dipholis salicifolia** A. de C., de C. Prod. viii, 188. **Xac-chum.**  
 "Tree 40 feet high, rare in the forests of Buena Vista Xbac, April" (1047). Flowers more densely crowded than the Florida U. S. form, and than 305 Flora Domingensis of Wright, Parry & Brummel.

## EBENACEÆ.

- Diospyros Ebenum** Kóen, Phys. Salsk. Hand. i, 176. **Tauch Ya.**  
*Sapote Negro.* Hacienda de Chablé (cult?) August 8, 1865 (*Schott, 542*).

## APOCYNACEÆ.

- Echites biflora** Jacq. Enum. Pl. Carib., 13.  
 In mangrove thickets at Sisal, Nov. 9, 1865 (*Schott, 812*).
- ECHITES SUBSAGITTATA Ruiz & Pav. Fl. Peru., ii, 19.  
*E. microcalyx* A. de C., Hacienda Saragossa, September 8, 1865 (*Schott, 674*).

**Echites torulosa** Linn. Sp. Pl. ii, 307.

"Vine 10 feet, common in brushlands about Izamal, producing its lemon-yellow flowers in August" (806, 883).

**Echites umbellata** Jacq. Enum. Pl. Carib., 13.

Sisal, May 9, 1865 (*Schott, 38*).

**Macrosiphonia Berlandieri** A. Gray, Syn. Fl. Am. ii, i, 83.

Yucatan, loc. ignot. et sine die (*Schott, sine num.*).

**Nerium Oleander** Linn. Sp. Pl. 305.

*Narciso.*

Yucatan loc. ign. (*Valdez, 8* "Colorada, e blanca"). Probably from cultivated plants at Merida. Merida. Cult. April, 1865 (*Schott, 813*).

**Plumeria pudica** Jacq. Enum. Pl. Carib., 13.

Cult? Camino de Sisal near Pozo del Toro, May 7, 1865 (*Schott, 428, 429*).

PLUMERIA RUBRA Linn. Sp. Pl., 209.

Cult? Merida, April to August, 1865 (*Schott, 321, 323, 324* flor rubro; 670 flor alba). All of these labels are associated with one specimen of the above species.

RAUWOLFIA HETEROPHYLLA Willd. ex. Roem. & Schult. Syst. iv, 805.

Suburbs of Merida, April 14, 1865 (*Schott, 431*).

**Stemmadenia insignis** Miers. Apocyn. S. Am., 76.

*Laurel.*

"Merida" (*Schott, type 430*). Merida, April, 1887 (*Millsbaugh, 27*). Only seen in cultivation in the public parks. The specimens in Herb. Schott bear a close resemblance to *Tabernaemontana citrifolia* Linn.

TABERNÆMONTANA ACAPULCENSIS Miers Apocyn. S. Am., 57. **Uçupek.**

"Merida, March, 1865 (*Schott, type 432*). Along fence rows, campo beyond Uman, April, 1887 (*Millsbaugh, 34*).

THEVETIA NEREIFOLIA Juss. ex. Steud. Nom. ed. 2, ii, 680.

Suburbs of Merida, April 11, 1865 (*Schott, 321*). Quinta del Obispo (*Schott, sine die sine num.*).

**Thevetia spathulata** sp. nov.

Foliis ovato-spathulatis vel (junior) oblongeolatis obtusis vel subacutis, margine integris, supra glabris et nitidis obsolete reticulato-venosis nervis lateralibus, centrali obliquus subtus pallidis glabris, cyma terminalia laxiflora glabra, bracteis triangulo-lanceolatis, deciduis, lobis calycinis lanceolatis mucronato-acuminatis glabris, tubo corollæ vix brevioribus. Frutex 20 pedalis Ramuli teretes glabris, cortex albo-chartaceis. Folia 2-4 poll. long., 6-14 lin. lat., petiolo 2-4 lin. long. Cymæ folio subsequantibus. Flores aurea-viridis 2 poll. long., pedicellis poll. long. Lobi calyces 5 lin. long., 2 lin. lat. Tubus corollæ intus apice strigose hirtis, appendicibus 5 cornatiformis. Affin. *T. cunifolia*.

Although there is a great variation in the shape of the juvenile leaves—the mature only being constant and typical—I have little hesitancy in referring the following young plants to this species, *Schott*, 321, Merida, 1865; *Gaumer*, 7, Cozumel Island, 1885 “new species?” (Oliver); and *Gaumer*, 678, “common at the Port of Silam, April, 1895.” Road beyond Xcholac, 15 feet high, flowers lemon-yellow, March 2, 1890 (*Stone*, 255, in *Herb. Acad. Sci.*, Phila.).

VINCA ROSEA Linn. Syst. ed x, 944.

Suburbs of Merida, common as a roadside weed. April 14, 1887 (*Millspaugh*, 44).

#### ASCLEPIADACEÆ.

ASCLEPIAS CURASSAVICA Linn. Sp. Pl., 314.

“Herb 3 feet high, abundant in damp situations near Izamal, its yellow flowers produced throughout the year” (556). Suburbs of Merida, January 5, 1865 (*Schott*, 182).

METASTELMA SCHLECHTENDALII Decne. in de C. Prod. viii, 513.  
Tekanto, Feb. 25, 1890 (*Stone*, 203, in *Acad. Sci.*, Phila.).

#### CONVOLVULACEÆ.

EVOLVULUS ALSINOIDES Linn. Sp. Pl. 392.

“Herb 18 inches high, common in the forests of Buena Vista Xbac, April” (1067). Merida, June, 1865 (*Schott*, 64).

IPOMŒA BONA-NOX Linn. Sp. Pl. ed. ii, 228.  
Nohcacab, Oct. 20, 1865 (*Schott*, 684).

**Ipomœa cissoides** Griseb. Fl. Brit. W. Ind. 473.

“Vine 10 ft., abundant in old fields about Izamal, producing its satiny-white flowers in November” (982). Range between Sacacab and Tikul, Oct. 20, 1865 (*Schott*, 905).

IPOMŒA COCCINEA Linn. Sp. Pl. 228.

“Vine 20 ft., common in old fields about Izamal, producing its bright red flowers from October to December” (981, 990).

IPOMŒA FASTIGIATA Sweet Hort. Brit. ed. i, 288.

“Vine 15 ft., common in old fields near Izamal” (915).

**Ipomœa hederacea** Jacq. Coll. i, 124.

“Vine 20 feet, common in towns and the fields about Izamal, blossoming in October and November. Flowers deep sky-blue changing to rose on drying” (919, 983). Merida, 1865. (*Schott*, 906).

“Vine 30 feet, rare at Buena Vista Xbac; flowers bright rose-pink” (769). In *Gaumer's* 919 and 983 the flowers are large and long peduncled and the leaves 7 inches broad by 5 inches long,



and scattered strigose-hairy. In Schott's 906 the leaves are small, about 2 by 1½ inches, strigose-hairy like the preceding. In Gaumer's 769 the leaves are small as in Schott's form, but are densely covered both above and beneath with long closely-apressed silky hairs.

***Ipomœa pentaphylla* (L.) Jacq. Coll. ii, 297. Soac**

"Vine 20 feet, abundant on roadsides about Izamal, producing its white flowers in October" (985). Merida, Nov. 7, 1865. (*Schott, 26*).

***IPOMŒA PUNCTICULATA* Benth. Bot. Voy. Sulph. 136.**

"Vine 20 feet, abundant in towns and fields, producing its reddish-purple flowers from September to February" (329).

***Ipomœa sinuata* Orteg. Hort. Matr. Dec. 84.**

"Vine 10 feet, frequent in brushlands and cornfields about Izamal" (1093)

***JACQUEMONTIA ABUTILOIDES* Benth. Voy. Sulph., 34.**

"Mountains between Hacienda Sacacal and Ticul, Oct. 20, 1865 (*Schott, 681*). Tekanto, Feb. 1890 (*Stone, 206*, in Herb. Acad. Sci. Phila.).

***JACQUEMONTIA PENTANTHA* G. Don., Gen. Syst. iv, 283. Yaxhal.**

"Vine 10 feet, common in brushlands about Izamal, October to January" (917). Suburbs of Merida, Nov. 14, 1864. (*Schott, 344*). Tunkas, March 3, 1890 (*Stone, 231, 235*, in Herb. Acad. Sci. Phila.).

***Jacquemontia tamnifolia* Griseb. Fl. Brit. W. Ind. 474.**

Vine 20 feet, common in the forests of Buena Vista Xbac, Aug. (1121).

· BORRAGINACEÆ.

***CORDIA DODECANDRA* de C. Prod. ix, 478.**

Thirty feet high, flowers vermilion with a tinge of orange; Tekanto, Feb. 27, 1890 (*Stone, 198*, in Herb. Acad. Sci. Phila.).

***CORDIA GERASCANTHOIDES* H. B. K. Nov. Gen. et. sp. iii, 69. Bohom.**

"Tree 100 feet high, common in forests where it blooms in April. Flowers very profuse, aromatic and durable. This tree yields one of the most valuable timbers of the country, much used in wagon-making and wherever strength and elasticity are required. Buena Vista Xbac" (1066). Aguada San Antonio, Xcanhacan, March 1, 1866 (*Schott, 745*).

***CORDIA GLOBOSA* (L.) H. B. K., Nov. Gen. et Sp. iii, 76.**

Merida, 1865 (*Schott, 970*). "Shrub 12 feet high, rare, Kantunil road near Izamal, producing its white flower in June" (745). "Shrub 30 feet high, in brushlands near Izamal, Sept." (801).

- CORDIA SEBESTENA** Linn. Sp. Pl., 190. **Kopté**  
Merida, Dec. 14, 1864 (*Schott, 173*); Sisal, May 10, 1865  
(*Schott, 386*).
- EHRETIA TINIFOLIA** Linn. Syst. ed. x., 936. **Roble.**  
Merida, Apr. 14, 1865 (*Schott, 800*).
- HELIOTROPIUM CURASSAVICUM** Linn. Sp. Pl., 130.  
Ticul, March 12, 1890 (*Stone, 275*, in Herb. Acad. Sci., Phila.).
- Heliotropium fruticosum** Linn. Syst. ed. x, 913.  
Merida, June, 1865 (*Schott, 364*).
- HELIOTROPIUM INUNDATUM** Sw. Prod. Veg. Ind. Occ., 40.  
Merida, Aug. 21, 1865 (*Schott, 968*).
- HELIOTROPIUM PARVIFLORUM** Linn. Mant. ii., 201.  
Merida, January 4, 1865 (*Schott, 175*).
- TOURNEFORTIA GNAPHALODES** R. Br. Prod., 496.  
Downs at Progreso, April 3, 1865 (*Schott, 305*).

## VERBENACEÆ.

- AVICENNIA NITIDA** Jacq. Enum. Pl. Carib., 25. **Mangle blanco.**  
Celestun, May 12, 1865 (*Schott, 361*).
- AVICENNIA OFFICINALIS** Linn. Sp. Pl., 110.  
*A. tomentosa* Jacq. Celestun, May 12, 1895 (*Schott, 473*).
- Citharexylum quadrangulare** Jacq. Enum. Pl. Carib., 26.  
Merida, Aug. and Sept. 1865 (*Schott, 89* et spec. *sine num.*)
- CLERODENDRON ACULEATUM** (L.) Griseb. Fl. W. I., 500.  
Merida, (*Schott, 27*).
- Clerodendron Colebrookianum** Walp. Rep. iv, 114.  
"Rarely cultivated, one bush 4 feet high at Izamal blooming  
in Sept. (909).
- DURANTA PLUMIERI** Jacq. Select. Am., 186.\* **Xcambocoché.**  
*D. repens* and *spinosa* L. Merida, June 11, 1865 (*Schott, 381*).
- LANTANA CAMARA** Linn. Sp. Pl., 874. *Oregano Silvestre.* **Xo-hexnuc.**  
"Shrubby, common in brushlands about Izamal, flowering  
from August until January" (808). "Abundant in a wild state"  
(*Valdez, 60*). Tekanto, shrub 6 ft. high, Feb. 25, 1890 (*Stone,*  
*213*, in Herb. Acad. Sci., Phila.).

\*Jacquin's name (1763) is accepted by Linnaeus in his third edition Sp. Pl. 1764.

**Lippia umbellata** Cav. Ic., ii., 75.

"Shrubby, 10 feet high, uncommon, found only on the ancient mounds southwest of Izamal" (871, 971). Our form is not so densely hairy either on the involucre or the under surface of the leaves as 4389, Heyde & Lux, Guatemala.

## PETREA ARBOREA H. B. K. Nov. Gen. et sp. ii., 282.

"Uncommon at Calcehtoh. A woody vine, with violet-like flowers deeper colored than the purple calyx, and with a spot on the middle petal." March 11, 1890 (*Stone*, 259, in Herb. Acad. Sci., Phila.).

## PRIVA LAPPULACEA (L.) Pers. Syn., ii., 1807.

**Oallunçay.**

*P. echinata* Juss. Suburbs of Merida, and Quinta del Obispo, Nov. to Dec., 1864 (*Schott*, 22 and 23). Tekanto, Feb. 27, 1890 (*Stone*, 186, in Herb. Acad. Sci., Phila.).

## LABIATÆ.

**Cedronella Mexicana** Benth. Lab. Gen. et Sp., 502. *Toronjil.*  
Campo near Merida, April 14, 1887 (*Millspaugh*, 45).

## HYPTIS PECTINATA (L.) Poit., Ann. Mus. Par., vii., 474.

"Herb 5 feet high, common on brushlands about Izamal, November" (943).

**Hyptis suaveolens** (L.) Poit., Ann. Mus. Par., vii., 472.

"Herb 3 feet high, common along roads and in old fields about Izamal, November" (942).

**Leonurus glaucescens** Bunge, Ledeb. Fl. Alt., ii., 409.  
Near Kikil, March 27, 1866 (*Schott*, 773). Cult?

## OCIMUM BASILICUM Linn. Sp. Pl., 833.

*Alvaca.*

Campo near Merida, escaped? April 14, 1887 (*Millspaugh*, 44).

## OCIMUM MICRANTHUM Willd. Enum. Hort. Berol., 630.

Suburbs and streets of Merida, June 23, 1865 (*Schott*, 511). Yucatan loc. ignot. (*Stone*, sine num., in Herb. Acad. Sci., Phila.). Calcehtoh, March 11, 1890 (*Stone*, 265, in Herb. Acad. Sci., Phila.).

**Rosmarinus officinalis** Linn. Sp. Pl., 33.

Campo at Uman, escaped, April, 1887 (*Millspaugh*, 31).

## SALVIA COCCINEA Buchoz. ex Etling. Salv., 23.

Merida, suburbs, April 14, 1865 (*Schott*, 160, loc. ignot., 760, in Herb. U. S. Natl. Mus.). Sitalpech, and Tekanto, Feb. and March, 1890 (*Stone*, 188, 256, in Herb. Acad. Sci., Phila.).

## SOLANACEÆ.

- Capsicum annum** Linn. Sp. Pl., 270. *Chile.*  
Cultivated at Merida, Feb'y 11, 1865 (*Schott, 237*). Campo  
near Merida, April 14, 1887 (*Millsbaugh, 43*).
- CAPSICUM FRUTESCENS** Linn. Sp. Pl., 189.  
Ticul (*Stone, 277*, in Herb. Acad. Sci., Phila.).
- CESTRUM DIURNUM** Linn. Sp. Pl. 277.  
Suburbs of Merida, April 28, 1865 (*Schott, 123*).
- DATURA STRAMONIUM** Linn. Sp. Pl., 255. *Chamico. Mehen-Xtocu.*  
"Herb 4 feet high. A very common weed in places, especially near small towns. In bloom from June to December, flowers pure white" (1062).
- Nicotiana glutinosa** Linn Sp. Pl., 259. **Kooo.**  
Campo near Merida, May, 1865 (*Schott, 486*).
- Nicotiana pilosa** Dunal., de C. Prod., xiii., i., 559. **Kooo.**  
"Waste grounds, campo near Uman, April, 1887 (*Millsbaugh, 39*).
- Physalis** sp.  
"Shrubby below, 3 feet high, very rare, found in fruit only, Sept. and Oct., along the road to Merida near Izamal" (1000).
- Physalis angulata** Linn. Sp. Pl., 262. *Jerba Mora.*  
"Herb 4 feet high, abundant in fields and gardens about Izamal, in blossom throughout the year" (479).
- Physalis foetens** Poir., Encycl. Suppl., ii., 348.  
Merida, Nov. 28, 1864 (*Schott, 5*).
- Physalis pubescens** Linn. Sp. Pl., 262.  
Suburbs of Merida, Dec. 4, 1864 (*Schott, 5*).
- SOLANUM AMAZONICUM** Ker-Gawl., Bot. Reg., t., 71. **Xkon-yaxik.**  
"Shrubby, 6 feet high, abundant in opens and brushlands about Izamal, producing its bright-blue flowers throughout the year" (366). Tunkas common, flowers bright purple," March 3, 1890 (*Stone, 234*, in Herb. Acad. Sci., Phila.).
- Solanum calicarpæfolium** Kth. and Bou., Ind. Sem. Hort. Berol., App. 10.  
Shrub 4 feet high, uncommon at the port of Oilam, producing its white flowers in April (652).
- Solanum fuscatum?** Linn. Sp. Pl. ed., ii., 268. **Xpacanul.**  
Merida, July 25, 1865; Cenoté Yopop, on the road from Merida to Progreso, April 7, 1865 (*Schott, 310*). Grisebach sp. cit. Flor. Brit. W. I. places Yucatan in brackets after this species without farther reference to authority.

**Solanum Jamaicense** Mill. Gard. Dict., Ed. viii. 17.

"Shrubby, 10 feet high, uncommon in brushland about Izamal, April to Aug." (595).

**Solanum Juripeba** Rich, Soc. Hist. Nat., Par. i. 107. **He-Bech.**

"Shrubby 10 feet high, common in the forests about Izamal, producing its greenish-white flowers from Sept. to Oct." (856). In our specimens the thorns are flatly compressed and straight, spinescent on the midrib above and below, also not rare on the calyx, the leaves are subentire and very unequal at the base.

**Solanum Mammosum** Linn. Sp. Pl., 267.

"Herb 10 feet high, frequent on roadsides and in forests about Izamal, blossoming in October" (998).

**Solanum Melongena** Linn. Sp. Pl., 266.

*Berengena.*

Cultivated at Merida (*Schott*, 238), not in flower Feb. 11, 1865.

**Solanum rostratum** Dun., Hist. Solan., 234.

"A common herb at Tekax, producing its lemon-yellow flowers in September" (1137).

SOLANUM TORVUM Sw. Prod. Veg. Ind. Occ., 47.

"Shrubby, 4 feet high, common at Progreso" (1167)

SOLANUM VERBASCIFOLIUM Linn. Sp. Pl., 263.

"Shrubby, 10 feet high, abundant in waste places and brush lands, flowering throughout the year, Izamal" (581). Suburbs of Merida, April 14, 1865 (*Schott*, 339).

SCHWENKIA AMERICANA Linn. Gen., ed. vi. 567.

"Along railroad near Izamal" (969 pt.).

#### SCROPHULARIACEÆ.

**Angelonia angustifolia** Benth., de C. Prod., x. 254. *Boca de la Viaja.*

"Herb 1 to 3 feet high, rarely cultivated at Izamal, where it produces its deep-blue flowers throughout the year" (544). Cultivated for its medicinal properties" (*Valdez*, 59).

BUCHNERA MEXICANA Hemsl. Biol. Centr. Am. Bot., ii. 457.

Grassy fields at Nohpat, Nov. 24, 1865 (*Schott*, 826).

CAPRARIA BIFLORA Linn. Sp. Pl., 628.

"Suffrutescent 4 feet high, common in brushlands about Izamal, March" (409, 590 in part). "Growing wild near Merida" (*Valdez*, 98). Merida, about houses, Nov. 30; and at the Cenote Jambula Dec. 14, 1864 (*Schott*, 455, 70).

**Conobea pusilla** (Benth.) Benth. & Hook. f. Gen., ii. 951.

"Herb 3-8 inches high, common in shady places near Izamal, producing its violet-blue flowers from July to August" (764).

CAPRARIA SAXIFRAGÆFOLIA Cham. & Schl. in Linn., v. 105.

Tekanto, flowers light blue, only one specimen seen, Feb. 28, 1890 (*Stone, 182*, in Herb. Acad. Sci., Phila.).

MONNIERA MONNIERA (L.) Britt., Mem. Torr. Club., v. 292. **Xaxcach.**

“Prostrate herb, abundant at the Aguada de Chulubmay, 9 miles east of Izamal, March” (1096).

MONNIERA PROCUMBENS (Mill) Kuntze Rev. Gen., 463.

*Herpestis chamaedryoides* H. B. K. Merida, Aug. 29, 1865 (*Schott, 616*). Tekanto, Feb. 27, 1890 (*Stone, 195*, in Herb. Acad. Sci.). Phila.

#### BIGNONIACEÆ.

**Adenocalymna** sp.

**Chacanicab.**

“A bijuco, very common in the forests about Izamal, producing its gamboge-yellow flowers in April and May” (575).

**Bignonia æquinoctialis** Linn. Sp. Pl., 869.

**Chacanicab.**

“Vine 40 feet, frequent in the forests near Buena Vista Xbac, April” (1068, 1098). Yucatan loc. ignot. (*Schott, 399*), “universally used as binding in constructing thatch roofs.”

**Bignonia diversifolia** H. B. K. ? Nov. Gen. iii., 133. **Chacxnetoloc.**

“Vine 40 (795) to 100 feet, uncommon in the forests about Izamal, producing its purple flowers from June to Sept.” (725, 795). **Jolak, Xcolak**, Merida, June 1 and Aug. 6, 1865 (*Schott, 406, 560*).

**Bignonia mollis** Vahl? Ecl. Am. ii., 46.

**Sacac.**

“Vine 40 feet, common in the forests about Izamal, producing its reddish-purple flowers in June” (739).

**Bignonia Unguis-cati** Linn. Sp. Pl. 623.

**Xcanlol-ak.**

“Vine 20 feet, common in the forests about Izamal, flowers yellow, produced in March” (507). Merida, May 20, 1866 (*Schott, 776*).

CRESCENTIA CUJETE Linn. Sp. Pl., 626.

**Huas.**

Merida, June, 1865 (*Schott, 521*).

**Parmentiera cereifera** Seem., Bot. Voy. Herald, 182. **Xcat-cunc.**

Merida, Aug. 25, 1865 (*Schott, 382*). Leaves, inclusive of the petiole, 1 inch long. “Shrub 12 feet high, common about Izamal, producing its neutral-green flowers throughout the year” (338). Leaves, including petiole, 2 to 2½ inches long.

PARMIENTERA EDULIS Mocq. de C. prod. ix., 244.

**Kaat.**

Merida, Quinta del Obispo, cult? March 12, 1865 (*Schott, 242*).

**Pithecoctenium hexagonum** de C. Prod. ix., 195. **Xnetoloc.**

"Vine 50 feet, common in the forests about Izamal, the creamy-white flowers produced in May and June" (702). Mr. J. H. Thompson sends me a fine example of the fruit, remarking that the Mayas use the seeds as a remedy for frontal headache, by moistening and applying them to the forehead or temples according to the location of the pain.

**Tabebuia** sp.**Xha-hau-ché.**

"Tree 30 feet high, rare in the forests of Buena Vista Xbac, April" (1048). Flowers only collected.

## TECOMA STANS Juss. Gen. 139.

Calcehtoh and Tunkas, bush 6-7 feet high, March 11, 1890 (*Stone*, 258, in *Herb. Acad. Sci. Phila.*).

## PEDALIACEÆ.

## MARTYNIA DIANDRA Glox. Obs. t. i.

At the Hacienda Saragossa, Sept. 8, 1865 (*Schott*, 623).

## ACANTHACEÆ.

## BOUCHEA PRISMATICA (Linn.) Kuntze Rev. Gen. 502.

"Herb 4 feet high, common at Yaxcaba and Tekax, October" (1139).

## BRAVAISIA TUBIFLORA Hemsl.

A common bush on the sand hills of Progreso, March 29, 1890 (*Stone*, 287, in *Herb. Acad. Sci., Phila.*).

DIAPEDIUM ASSURGENS (Linn.) Kuntze, Rev. Gen., 485. *Pensamiento.*

Suburbs of Merida, Dec., 1864 (*Schott*, 154).

## HENRYA COSTATA A. Gray.

*Tetramerium costatum.* Tunkas, March 3, 1890; 2 feet high, flowers yellow striped above with reddish brown (*Stone*, 244, in *Herb. Acad. Sci., Phila.*).

## RUELLIA ALBICAULIS Bert. Spreng. Syst., ii., 822.

Merida, Nov. 1864 (*Schott*, 2).

**Ruellia geminiflora** H. B. K. Nov. Gen. et. Sp. ii., 240.

Tekanto, flowers lilac, only one plant seen, Feb. 27, 1890; Tunkas common, March 3, 1890 (*Stone*, 185, 242, 247, in *Herb. Acad. Sci. Phila.*). Dr. Gaumer's 384, collected Oct. 1895 near Izamal, and referred on page 320 doubtfully to *R. Tweediana*, is doubtless also this species.

## RUELLIA TUBEROSA Linn. Sp. Pl., 635.

**Cabal-Yaxnic.**

Merida at Quinta del Obispo, Nov. 24, 1864, and April 13, 1865 (*Schott*, 45, 282). Izamal, Feb. 28, 1890 (*Stone*, 254, in *Herb. Acad. Sci. Phila.*).

**SIPHONOGLOSSA SESSILIS** (Jacq) Oerst.

“Suffrutescent 2 feet high, common in waste places near Izamal, April” (532).

**TUBIFLORA SQUAMMOSA** (Jacq) Kuntze, Rev. Gen., 500.

Merida, at Quinto del Obispo, Nov. 23, 1864 (*Schott*, 24).

**VALERIANODES JAMAICENSE** (Linn.) Boerh. (Med.) Phil. Bot. i., 177.

“Herb 4 feet high, common at Merida, Izamal and inland” (1160).

## PLANTAGINACEÆ.

**PLANTAGO MAJOR** Linn. Sp. Pl., 163.*Llantén.*

Merida, at Quinta del Obispo, Jan. 18, 1865 (*Schott*, 193).

## RUBIACEÆ.

**ERNODEA LITTORALIS** Sw. Prod. Veg. Ind. Occ. 29.

Downs at Progreso, April 4, 1865 (*Schott*, 287).

**Exostemma Caribæum** (Jacq.) R. & S. Syst. v. 18. **Sabac-ché.**

“Shrub 20 feet high, not common at Xcholac” (957). Merida, Aug. 20, 1865 (*Schott*, 585).

**CHIOCOCCA RACEMOSA** Linn. Sp. Pl. 246.**Xcanchac-ché.**

Merida, July 28, 1865 (*Schott*, 569, 499).

**Coffea Arabica** Linn. Sp. Pl., 245.*Café.*

Cult? Merida, in flower and fruit Jan. 1866 (*Schott*, *sine num*).

**CRUSEA CALOCEPHALA** de C. Prod. iv., 567.

“At the Hacienda Mucuyché on road from Merida to Sacalum” (*Schott*, 676). “Herb 3 feet high, common on old fields, and in waste town lots at Izamal” (937).

**HAMELIA PATENS** Linn. Sp. Pl., 246.**Kanan.**

Suburbs of Merida, and in gardens Nov. 30, 1864 (*Schott*, 174).

**MORINDA ROYOC** Linn. Sp. Pl., 250.**Hoyoc.**

Near Merida (*Valdez*, 67). At Nohcacab, Nov. 27, 1865 (*Schott*, 709). Tunkas, a shrubby climber, March 3, 1890 (*Stone*, 240, in Herb. Acad. Sci. Phila.).

**Psychotria undata** Jacq. Hort. Schoen. iii., 5.*Cancerillo.*

“The dried and powdered, or toasted leaves used in the cure of **Uaga**, a form of malignant pustule.” Merida, July 6, 1865 (*Schott*, 524). Differs from the description only in having some of the stipules awn-pointed.

**RANDIA ACULEATA** Linn. Sp. Pl., 214.

North of Merida on road to Progreso, April 7, 1865 (*Schott*, 262). Merida (*Schott*, *sine num* in Herb. U. S. Natl. Mus.).



SPERMACOCE TENUIOR Linn. Sp. Pl. 147.

“Herb 18 inches high, frequent in shady situations near Izamal” (965). Merida, at Quinta del Obispo, Nov. 24, 1864 (*Schott, 57*). Nohcacab, Oct. 19, 1865 (*Schott, 972*).

SPERMACOCE VERTICILLATA Linn. Sp. Pl., 148.

Merida, 1865; Nohcacab, Oct. 19, 1865; Ruins of Uxmal, Sept. 16, 1865 (*Schott, 117, 971, 972*). Tekanto common, Feb. 25, 1890 (*Stone, 207*, in Herb. Acad. Sci., Phila.).

#### CAPRIFOLIACEÆ.

SAMBUCUS CANADENSIS Linn. Sp. Pl., 269.

Merida, (Cult?), Jan. 3, 1865 (*Schott, 180*).

#### CUCURBITACEÆ.

CAYAPONIA ALATA Cogn. de C. Monogr. 3.

Hacienda Saragossa, Sept. 8, 1865 (*Schott, type 901*).

CAYAPONIA GRANDIFLORA Cogn., de C. Monogr. 3.

Ticul, Oct. 21, 1865 (*Schott, type 680*).

**Lonicera macrantha** ? (Don.) de C. Prod. iv. 333.

Vine 10 feet, uncommon in the forests of Tekax, September (1105).

LUFFA ÆGYPTIACA Mill., Gard. Dict., ed. 8.

*L. cylindrica* Rœm. Suburbs of Merida, November 30, 1864 (*Schott, 6*).

MELOTHRIA PENDULA Linn. Sp. Pl., 35.

Merida, June, 1865 (*Schott, 572*). “The leaves are used by the Mayas to extract foreign bodies from beneath the eyelids.” Leaves less lobed than in other Yucatan collections.

CORALLOCARPUS MILLSPAUGHII Cogn. F. Col. Mus. Bot., i, 322. **Xlucioan.**

Merida, July 11, 1865 (*Schott, 479*).

**Pittiera longipedunculata** Cogn., Bull. Soc. Bot. Belg. xxx, 272.

**Ponponcit.**

“A vine 10 feet, rare in moist places near Tekax, September” (1135).

**Cucumis Anguria** Linn. Sp. Pl. 1436. **U-habaplat-oil-Kax.**

“An infrequent vine, Tekax, September” (1127). “Vine 20 feet, rare, Progreso” (1158).

MELOTHRIA PENDULA Linn. Sp. Pl., 49.

“Vine 15 feet, common near Progreso” (1180).

**Melothria fluminensis** Gardn. Jour. Bot., i, 173.

“Vine 10 feet, uncommon near Progreso” (1181). Merida, November, 1865 (*Schott, 900*).

MOMORDICA CHARANTIA Linn. Sp. Pl. 1433.

"Vine 20 feet, common in waste places at Tekax, September" (1124). Hacienda Guayalcé, September 19, 1865 (*Schott, 633*).

TRICERATIA BRYONIOIDES A. Rich., Ess. Fl. Cub., 614.

*Scydlum tamnifolium* Cogn. "Vine 25 feet, common in stony soil and on stone fences near Izamal, October and November" (922).

#### GOODENIACEÆ.

SCAEVOLA PLUMIERI (L.) Vahl. Symb. ii, 36.

Sandy seashore at Progreso, April 5, 1865 (*Schott, 294*).

#### COMPOSITAE.\*

**Achillea Millefolium** Linn. Sp. Pl., 899.

*Alcanfor.*

Campo about Merida, not in flower, April, 1887 (*Millspaugh, 47*).

AGERATUM CONYZOIDES Linn. Sp. Pl., 839.

Near Merida, January 4, 1865 (*Schott, 208*).

ALOMIA AGERATOIDES H. B. & K., Nov. Gen. et. Sp. iv. 151.

Tekanto and Tunkas, February and March, 1890 (*Stone, 190, 238*, in *Herb. Acad. Sci., Phila.*).

BIDENS LEUCANTHA Willd Sp. Pl., iii. 1719.

"Herb 2 feet high, abundant at the Port of Silam, rays white, April" (632).

**Bidens tereticaulis** de C. Prod., v. 598.

Merida, December, 1864 (*Schott, 145*). Tunkas, March 3, 1890 (*Stone, 240*<sup>a</sup> in *Herb. Acad. Sci., Phila.*). "Ascending among shrubbery like a vine, producing its orange-yellow flowers in October" (951).

**Brickellia diffusa** A. Gray, Pl. Wright., i. 86.

"Herb 10 feet high, uncommon in fields and brushlands near Izamal, flowering from December to March" (1080).

**Calea axillaris urticifolia** Rob & Greenm. Proc. Am. Acad. Sci., xxxii, 27.

Shrubby 5-6 feet high, flowers yellow, Tunkas, March 3, 1890 (*Stone, 243, 245*, in *Herb. Acad. Sci., Phila.*). "Shrubby 15 feet high, ascending among shrubs in the brushlands near Izamal" (956).

**Calea Zacatechichi** Schl. in Linn. ix, 589.

**Xicin.**

Nohpat, November 24, 1865 (*Schott, 912*).

**Cosmos caudatus** H. B. & K. Nov. Gen. et. Sp. iv. 240. **Chacxul.**

"Herb 4 feet high, uncommon at Izamal" (940).

\* Determined (with the exception of *Pectis*) by Mr. J. M. Greenman.

DYSODIA CANCELLATA A. Gray, Proc. Am. Acad. xix, 38.  
Merida, December 4, 1864 (*Schott*, 78).

**Eclipta alba** Hassk. Pl. Jav. Rar., 528.

"Herb 10 inches high, in shady places near Izamal, uncommon, October" (1012).

ERIGERON CANADENSIS Linn. Sp. Pl., 863.

Charcos at Celestun, May 12, 1865 (*Schott*, 362).

EUPATORIUM CONYZOIDES Vahl. Symb. Bot., iii. 96.

Suburbs of Merida, December, 1864 (*Schott*, 144). "Ascending 15 feet, among other shrubbery and on old fences in the brushlands about Izamal, producing its heliotrope-colored flowers in October" (914, 953).

EUPATORIUM PYCNOCEPHALUM Less. in Linn. vi. 404.

"Herb 6 feet high, common at Aguadas near Izamal, October" (1022).

FLAVERIA LINEARIS Lag. Gen. et. Sp. Nov., 33, 407.

Celestun, May 12, 1865 (*Schott*, 408).

**Flaveria repanda** Lag. idem.

Downs of Progreso, December, 1865 (*Schott*, 973). "Merida and Progreso, along the coast" (1147 part).

**Helianthus annuus** Linn. Sp. Pl., 1276.

*Girasol.*

Campo, suburbs of Uman, escaped, April, 1887 (*Millspaugh*, 32).

**Isocarpha oppositifolia** R. Br., Tr. Lin. Soc., iii., 110.

Tizimin, March 28, 1866 (*Schott sine num.*). "Herb 5 feet high among the shrubbery of the brushlands and forests near Izamal, October to January" (976).

MELANTHERA HASTATA Michx., Fl. Bor. Am., ii., 106.

*Amellus niveus* (L.) O. K. Merida, May 30, 1865 (*Schott*, 164.)

MELAMPODIUM PALUDOSUM H. B. & K., Nov. Gen. et. Sp. iv., 273.

*M. divaricatum* de C. Merida, Nov. 14, 1864 (*Schott*, 28). Calcehtoh, Mar. 11, 1890 (*Stone* 272, in Herb. Acad. Sci. Phila.) "Common in open places near Izamal, Nov." (961).

**Mikania cordifolia** Willd., Sp., Pl., iii., 1746.

About Nohcacab, Nov. 1865 (*Schott*, 716).

**Millieria quinqueflora** Linn. Sp. Pl. 919.

**Xyoutoloc.**

Cercanias about Merida, Nov. 15, 1864 (*Schott*, 90). "Herb 9 feet high, abundant in fields and vacant lots about Izamal. Flowering from Oct. to Jan'y" (949).

**Montanoa grandiflora** Hemsl.? Bio. Cent. Am. Bot., ii., 165.

Tizimin, Mar. 28, 1866 (*Schott, sine num.*).

- Pectis elongata Schottii** Fernald.\* Proc. A. A. A. S., xxxiii., 78.  
 "Much more slender in its diffuse branches than the species, leaves small, heads scattered and solitary in the axils, on capillary 3-4-bracteate peduncles 2 or 3 cm long: involucre pale, 4 mm high; pappus hardly equaling the akene. Serceania near Maxcanú, Sept. 14, 1866" (*Schott, 666*, Cat. No. of Type 40483 Herb. Field Col. Mus.).
- Pectis linifolia** Linn. Syst., Ed. x. 1221.  
*P. pectinata* Jacq. "Herb 2 feet high, stone walls near Izamal, uncommon" (979).
- PECTIS PROSTRATA Cav., Ic., iv., 12.  
 "Herb 6 inches high, common along roads and paths in the forests about Izamal" (1092). Merida, Aug. 20, 1865 (*Schott, 533*).
- Plúchea camphorata** de C., Prod. v., 452.  
 "Herb 6 feet high, abundant at Chulubmay, producing its purple flowers in March" (1099).
- PLUCHEA ODORATA Cass. Dict. Sc. Nat. xlii., 3 **Chalché**  
 Salt swamps at Sisal, Nov. 9, 1865 (*Schott, 915*). Specimens gathered by Dr. Schott at Merida (185) and by myself at Izamal (*Armour Expd 176*) were doubtless introduced from the coast, as the plant is used in Yucatec domestic medicine much as is bone-set in the United States.
- PLUCHEA PURPURASCENS de C., Prod. v., 452.  
 "Herb 6 feet high, old fields about Izamal, uncommon, producing its rose-colored flowers in October" (1002).
- Porophyllum Ervendbergii** A. Gray, Proc. Am. Acad., xix., 35. **Xpeehukil.**  
 Merida, at the Quinta del Obispo, Nov. 15, 1864 (*Schott, 81*).
- SANVITALIA PROCUMBENS Lam., Jour. His. Nat. Par., ii., 176. **Xkantumbub**  
 Merida, where it is used by the Mestizas as a remedy for the spitting of blood, Nov. 29, 1864 (*Schott, 14*). "Herb 18 inches, common at Cacalchen and Euan, producing its yellow and red-brown flowers from June to March" (964).
- Sclerocarpus divaricatus** B. & Hook. f., Biol. Cent. Am. Bot. ii., 164.  
 Labcah, July 30, 1865 (*Schott, 67*). "Herb 3 feet high, abundant in old fields near Izamal, from July to Jan'y" (410 part, 771).
- Senecio Berlandieri** (de C.) Hemsl. Biol. Cent. Am. Bot., ii., 236.  
*Gynoxis* de C. A climber among mimosa trees near Sisal (*Schott, 407*).

\* The species of this genus were determined by Mr. M. L. Fernald.

**SYNEDRELLA NODIFLORA** Gaertn. Fruct. ii. 456.

Merida, at the Quinta del Obispo, Nov. 15, 1864 (*Schott, 93*).  
 "Herb 6-18 inches high, common in waste places near Izamal,  
 in bloom throughout the year" (359, 952).

**TAGETES PATULA** Linn. Sp. Pl., 887.

On the Teocali of Kabah, Nov. 26, 1865 (*Schott, 708*). Calcehtoh, Mar. 11, 1890 (*Stone, 273*, in Herb. Acad. Sci. Phila.).

**TITHONIA DIVERSIFOLIA** A. Gray, Proc. Am. Acad., xix. 5. *Arnica*.

"Herb 15 feet high, quite shrubby, cultivated at Izamal where it blooms in November" (944).

**Tithonia tagetiflora** Desv. Ann. Mus. Par., i., 46. **Zum**

Merida, Dec. 3, 1864 (*Schott, 68*). "Herb 10 feet high, abundant in old fields, producing its orange-yellow flowers from Aug. to Sept." (899).

**TRIDAX PROCUMBENS CANESCENS** de C., Prod. v., 679.

Tekanto, Feb. 27, 1890 (*Stone, 212*, in Herb. Acad. Sci. Phila.).

**TRIXIS FRUTESCENS** P. Br., Pl. Jam. 312.

*T. radialis* (L.) O. K. Merida, Feb. 2, 1865 (*Schott, 189*).

**VIGUIERA HELIANTHOIDES** H. B. & K. Nov. Gen. et. Sp., iv., 226. **Toh**.


Suburbs of Merida, Jan. 1, 1865 (*Schott, 190*). "Herb 2 to 5 feet high, very abundant in old fields about Izamal and Xcholac, Oct. to May" (502, 557).

**Xanthium Canadense** Mill. Gard. Dict., ed. viii., 2.

"Herb 2 feet high, uncommon near Merida where it has probably been introduced in foreign baled hay" (1145).

**Zexmenia Costaricensis** Bth. Oerst. Kjob. Medd. 95. **Sactah**.

Between Guayalceh and Yumuó, Oct. 2, 1865 (*Schott, 914*).  
 "A shrubby climber, 20 feet, on an artificial mound 8 miles southwest of Izamal, Nov." (960, 962).



Notes and New Species—Prof. Ludwig Radlkofer; A New  
Myginda—Dr. Theodor Loesener; and Additional  
Notes and Species by the Author.

ULMACEÆ.

**Mertensia lævigata** H. B. K. Nov. Gen. et. Sp., ii., 31.

*Celtis aculeata* Sw. ex Ind. Kewens., sed vix recte ob folia integerrima; inde *Celtis lævigata* nominanda—Radlkofer.

“Common at Buena Vista Xbac, flowering in May” (567).

CAPPARIDACEÆ.

**Forchhammeria trifoliata** Radlk. sp. nov. *Tres Marias.*

Arbor glabra; rami florigeri pallidi, lenticellosi, annulo ligni (librique) dupliori insignes; folia ad ramorum apices conferta, petiolata, trifoliolata; foliola ex obovato cuneata, breviter acutiuscule acuminata vel obtusata, breviter petiolulata, margine integerrimo subundulata, pallide viridia, fibris sclerenchymaticis crebris percursa; flores parviosepali, apetalii, masc. in paniculam amplam congesti, staminibus 12–15 disco insertis, fem. subracemosi, germine intra staminodia pauca disco insidente ellipsoideo 2–(rarius 3–) loculari, stigmatibus sessilibus 2(–3)–lobo, gemmulis in loculis binis ad media septa crassiuscule carnosa vel ad angulos centrales collateraliter affixis pendulis anatropis epitropis; fructus ignotus.

“*Arbor* 30-pedalis sat rara” (Gaumer). *Rami* florigeri diametro 6 mm. teretes, lenticellis crebris breviter ellipticis sulco mediano notatis obsiti, jam annulum secundum fasciculorum fibrovasalium ut in *Roydsia* et *Maerua* (cf. Radlkofer in Sitzungsber. K. bayer. Acad. XIV, 1884, p. 90) intra libri primarii fibros sclerenchymaticos (nec ut in *Forchhammeria pallida* Liebm. et F. apiocarpa Radlk. extra illos, cf. l. c.) ortum exhibentes. *Folia* innovationum inferiora sub gemmis dormientibus abortiva vel in squamulas conversa, superiora petiolo 3–10 cm. longo adjecto 14–24 cm. longa, 10–16 cm. lata. foliola petiolulatis 3–4 mm. longis adjectis 10–16 cm. longa, 3,5–6 cm. lata. lateralia paulominora subcoriacea, penninervia, nervo mediano supra planiusculo subto convexo sat valido siccitate undulato in apicem callosum excurrente, nervis lateralibus sat numerosis (12–18) dentibus oblique patulis ante marginem arcuato-anastomosantibus, inter nervos retivenarum obliquo (quasi vena ramificata ab arcu

ad nervum medianum recurrente) notata margine linea cartilaginea circumvallata, glaberrima, utrinque opaca vel supra subnitida, epidermide nunc conspicue, nunc vix crystallophora, diachymate arcte fibroso, hypodermate ad paginam superiorem vix nisi juxta nervos instructa; petiolus a dorso adventrem compressiusculus, striatus, supra sulco angusto notatus, basi incrassatus cum ramulo articulatus; stipulae minimae, apiculos glanduliformes ad mediam petioli insertionem (vel folii cicatricem) exhibentes. *Florum masc.* paniculae densiflorae, pyramidales, 17 cm. longae, 6 cm. latae, ramis racemiformibus praeter florem terminalem saepius luxuriantem flores complures laterales pedicellatos hic inde in dichasia consociatos in axillis bractearum parvularum subulatarum gerentibus, adjectis interdum floribus rudimentaritis; pedicelli 3-6 mm. longi. *Sepala* bracteis conformia, parva, 1.5 mm. longa, subulata, 4-6; *Petala* o. *Discus* pulvinatus. *Stamina* supra discum inserta, filiformia, glabra, antheris brevibus, basi subcordatis, subintrorsis, ad connectivi basin dilatata dorso insertis. *Pistilli* rudimentum o. *Florum foem.* inflorescentiae racemiformes (polychasia) in axillis foliorum 5 cm. longae, ima basi utrinque ramum aequilongum emittentur praeter florem terminalem, flores complures laterales, pedicellatos vel eorum loco hic inde dichasia vel polychasia parvula in axillis bractearum subulatarum vix 2 mm. longarum gerentes; pedicelli 2-3 mm. longi. *Sepali* bracteis conformia, 4-6. *Petala* o. *Discus* elevato-pulvinatus, carnosus. *Staminodia* 5-6, 1-seriata, stamina rudimentaria abbreviata antheris sterilibus exhibentia. *Germen* ellipsoideum, 2 mm. longum, 1.5 mm. latum, basi coarctata subsessile, plerumque biloculare; stigmatis lobi dorsales.

Yucatan: Dr. G. F. Gaumer n. 370! (flor. foem. m. Feb. 1895); n. 417! (flor. masc. m. Mart. 1895).\*

Obs. A Forchhammeriae specibus reliquis (*F. pallida* Liebm., *F. apiocarpa* Radlk. *F. Watsoni* Rose) recedit foliis compositis et floribus masculis paniculatis nec non foliorum fabrica (hypodermate minus conspicuo, epidermate interdum vix crystallophora, diachymate fibris percurso), insuper ligni annulo secundo intra libri primarii fibros (nec extra eos) oriente. Attamen vix dubie ejusdem generis est.

## MALPIGHIACEÆ.

### **Malpighia glabra acuminata** A. Juss.

Ex descr.—Radlkofer. "Shrub 15 feet high, not common in the forest and brushlands about Izamal, where it produces its showy rose-pink flowers in November" (972).

\*Branch of female shrub 20 feet high, collected at the base of the smaller mound at Izamal, January (*Millspaugh, A. V. Armour Exped.* 86).



## RUTACEÆ.

**Casimiroa tetrameria** Millsp. sp. nov.

Arbor glabra, 30 pedalis, rami pallidi verrucosi, foliis alterna terminalia longe (5-9 cm.) petiotata, dense et molliter hirsuta, digitatim 3-5-foliolata, foliolis ovato-lanceolatis integerrimis e basi cuneatis in acumen emarginatum terminatis, omnia breviter petioulata, duo inferiora frequenter auriculo-lobulata. Florum masc. in paniculae ramosae axillares et extra-axillares ad cicatricis foliorum antiquus, 2-7 cm. long. 1-5 cm. lat., pubescentibus, ramis plus minus racemiformibus; pedicellis floribus aequantis, bractæ minutæ stipuliformis, calyce tenuis 4-dentibus, petala 4, crassiuscula carnosa, stamina 4, erecta, infra discum inserta ad basi crassiuscula, antheris verticillatis, discus minutis. Flor. foem. et fruct. ignot.

Folia 18-24 cm. long., foliola 6-10 cm. long., 2.5-5 cm. lat. Rara ad Xcholac, m. Nov. (1006).

**Esenbeckia pentaphylla** Griseb. Fl. Brit. W.I., 135. **Yax-hocob.**

Ex descr.—Radlkofer. "Tree 50 feet high, uncommon in the forests about Izamal, producing its cream-colored flowers in July" (752).

## EUPHORBIACEÆ.

**Adelia Oaxacana** (Muell.) Hemsl. Biol. Cent. Am. Bot., iii. 129?**Xtompac.**

Ex descr.—Radlkofer. "An abundant shrub 20 feet high, Xcholac, February" (446).

## CELASTRACEÆ.

**Myginda Gaumeri** Loes. sp. nov.

Ramulis parva et tenuiter hirtellis; foliis progenere majusculis, ovata-lanceolatis, minute serrulatis, basi cuneatis, apice obtusis et minute apiculatis, tenuiter coriaceis, glabris, 5-9.5 cm. longis, 2-3.4 cm. latis, supra et subtus dense prominulo et reticulatim nervosis; inflorescentiis in foliorum axillis solitariis, dichotome furcatis, sub lente parce et tenuiter hirtis, bracteis minutis; floribus numerosis, 4-meris; calyce extrinsecus tenuissime pilosulo, sepalis obtusis; petalis rotundatis; disco lobis 4, subemarginatis formato; staminibus inter eius lobos insertis, parvis; ovario 2-loculari, in stylum brevissimum angustato; loculis 1-ovulatis, ovulis erectis, stigmatem parvo capitellatis.

Yucatan, Buena Vista Xbac: Gaumer n. 1049. Flor. m. Apr. Petala in sicco brunneo-violacea sed postquam in aqua cocta colorem glauco-coeruleum ostendunt. "Frutex sylvicola, 15 pedalis."

## HIPPOCRATEACEÆ.

- Hippocratea Grisebachii** Loes. Eng, Prantl. Nat. Pfl. iii.5, 227.  
*H. verrucosa* Griseb. in Goett. Abh. xxiv: 84 (1879) Non H. B.  
 K. Nov. Gen. et Sp., v. 136 (1821), teste Loesener ipso in litl.  
 "Vine 25 feet, common in forests about Izamal, February" (385).

## SAPINDACEÆ\*

- Allophylus Cominia** Swtz. Prod. Veg. Ind. Occ., 62 (1788).  
*Schmidelia Cominia* Swtz. Prod. Veg. Ind. Occ., ii. (1800).  
 "Shrub 20 feet high, common in forests about Izamal, August"  
 (893).

[CARDIOSPERMUM CORINDUM molle Radlk.

*C. molle* H. B. K.—*C. Corindum forma 3 molle*. Radlk. Flor.  
 Brazil Fasc. 122:447 (Sept. 1897.) Gaumer's 494 (Contrib.  
 ii:306) has not been seen by Prof. Radlkofer, the above note is  
 simply bibliographic.]

**Paullinia Cururu** Linn. Sp. Pl. 365.

"Vine 50 feet, common in the forests of Buena Vista Xbac,  
 April" (1041).

PAULLINIA FUSCESCENS GLABRESCENS Radlk.

*Paullinia Curassavica* of Biol. Centr. Am., Bot. iv:3, Mills-  
 paugh Contrib. Fl. Yuc. I:33, is to be replaced by *Paullinia fus-*  
*cescens* Kth., forma 3. *glabrescens* Radlk., Monogr. Paullin.  
 (1895-96) in Abhan. K. bayer. Acad., 283, (Sep. 217) line 10 from  
 below. The true *Paullinia Curassavica* of Linné, is a *Serjania*  
 -*S. Curassavica* Radlk., in Monogr. Paull., 305 (Sep. 239). The  
 plant collected on the Island of Cozumel by Gaumer and noted  
 in Biol. Cent. Am., Bot. iv:3 has been overseen (sic) in Radlk.  
 Monogr. Paull. The quotation of Grisebach Fl. Brit. W. Ind.  
 by Hemsley is right for the Columbian plant, not that of  
 Jamaica, which is *P. Jamaicensis* Macf.

"Vine 30 feet, common in forests about Izamal, Sept. and  
 Oct." (900).

**Sapindus Saponaria inaequalis** (de C) Radlk.

*Sapindus Saponaria* L. forma *inaequalis* Radlk., *S. inaequalis*  
 de C. This is the "*Sapindus* sp. near *divaricatus* Willd" pub-  
 lished in Biol. Centr. Am., Bot. iv. 3, collected by Dr. Gaumer  
 on the Island of Cozumel in 1886, which I have seen; also 396  
 Gaumer "common about Izamal, flowering in February."

(I observed this form growing plentifully at the Laguna  
 Columbia and Cedral, Island of Cozumel in 1895 while accompa-  
 nying the Allison V. Armour expedition, and collected it also at  
 Izamal. C. F. M.)

\*Professor L. Radlkofer.

**Serjania adiantoides** Radlk. sp. nov.**Bui.**

Fructu et habitu in Sect. xii. (*Synococcus*) affinis *Serjaniae* orbiculari Radlk., insignis caulibus subherbaceis in angulis pilis patentibus hirsutis, foliis sub-bipinnatis 4-jugis, jugo infimo utrinque 5-foliato pinnato, sequenti ternato, reliquis simplicibus, floribus in thyrsis bicirrosis ob rhachim abbreviatam subumbellatim confertis. (Habitu quo dammodo in mentem revocat *Pauliniam thalictrifoliam* Juss.).

Yucatan: Dr. G. F. Gaumer n. 1114 (Buena Vista Xbac, m. Sept. 1895, alab. et fruct.; stirps rara, ad altitudinem "10 ped." scandens.

**SERJANIA ATROLINEATA** Sauv. & Wr. Fl. Cuba, 24.

*Serjania Scatans* Radlk. Monogr. Serj. 213., see Radlk. Serj. Suppl., 1886:116. This replaces the *Seriana scatans* (Radlk.) on p. 33 of these contributions.

**Serjania goniocarpa** Radlk. Serj. Monogr. 309.

"Vine 60 feet, abundant in forest and brush lands about Izamal where it flowers in September" (973).

**SERJANIA PLICATA** Radlk. Monogr. Serj. 167.

Not *Seriania plicata* (Radlk.) as listed on p. 33 of these contributions.

**Talisia olivaeformis** (Kth.) Radlk., in Sitz. K. bayer. Acad., viii. 342.*Guayo. Uayum.*

*Melicocca olivaeformis* Kunth. "Tree 60 feet high, common in the forests of the peninsula, and cultivated in the villages. February to April" (406).

**Thouinia paucidentata** Radlk. sp. nov.

Arbor "30 pedalis, vulgaris" (Gaumer); folia ternata (petiolo 2 cm. longo adjecto 6-7 cm. longo); foliola lanceolata, in acumen obtusum terminata, supra medium remote obtuso-dentata, praeter nervorum axillas subtus barbatae glabra, impunctata, epiderme mucigera; thyrsi axillares, racemiformes, eramosi, breves, stricti, puberuli; flores in axillis bractearum per brevium plerumque bini, pedicellati, puberuli. (Fructus non suppetebant.)

Yucatan: Dr. G. F. Gaumer n. 865! (m. Aug.-Sept. 1895, flor.)

**Urvillea ulmacea genuina** Radlk.

*Urvillea ulmacea* Kunth, forma 1 genuina Radlk., in Sitz. K. bayer. Acad., viii. (1878): 263, 264; et in Fl. Braz. Fasc. 122 (1897): 423.

"Vine 20 feet, abundant in brush lands about Izamal, January to July 481."

(Through an error in the printer's make-up of p. 307, Contribution ii., this species is included under *Vitaceae* instead of preceding that order at the head of the pagé.)

## VIOLACEÆ.

**Hybanthus Yucatanensis** Millsp. sp. nov.

Frutex "8 pedalis," caule albo erecto, sparsim aculeato, ramoso; ramis junior. viridis quadrangularis, angulis tenuiter papyraceo-alatis; foliis ramibus plerumque fasciculatis vel alternis, cuneato-lanceolatis margine regulariter crenato-dentatis, apice plerumque acutis, quandam obtusis rarissime emarginatis, petiolo plus minus patentibus, rarissime 1-2 mm. longo. Floribus minimis fasciculatis pedunculatis; pedicellis unifloris farinaceis. Sepalis ovatis acutis margine ciliatis, petalo infimo calyce duplo longiore, basi saccato apice cucullato integerrimis, lateralibus et superioribus ovatis obtusis margine erosis; antheris columnaris, glandula nectarea magna conchæformis, stylo nepenthæformis, ovario ovato subtrisolcato (junior.).

"Infrequens ad Izamal, Yucatan m. Feb.-Mart. n. 469, flores virido-flava." Dr. G. F. Gaumer, 1895.

## SAPOTACEÆ.

**Bumelia obtusifolia buxifolia** Miq. et Eichl. Fl. Bras. vii: 47.**Puomucuy.**

"Shrub 15 feet high, abundant in brushlands about Izamal, March to April" (473).

# Index.

	PAGE.		PAGE.
Abutilon .....	377	Arisæma .....	355
Acacia.....	363	Aristida .....	350
Acalypha .....	370	<i>Aristida</i> .....	350, 354
ACANTHACEÆ .....	391	Aristolochia .....	359
Achillea.....	394	ARISTOLOCHIACEÆ.....	359
Achras .....	382	Artocarpus .....	358
Acrocomia .....	354	Arum .....	355
Adelia .....	400	<i>Arundinaria</i> .....	354
Adenocalymna .....	390	Arundo .....	350
Adiantum .....	348	ASCLEPIADACEÆ.....	384
Æchmea .....	355	Asclepias .....	384
Æschynomene .....	363	Aspidium .....	348
Ageratum .....	394	Asplenium .....	349
ALGÆ.....	347	Astronium.....	375
Alicastrum.....	358	Atriplex.....	359
Allionia .....	361	Attalea .....	355
Allophylus .....	402	Avicennia .....	386
Alomia.....	394	Ayenia .....	379
Alternanthera .....	360		
Alvaradoa.....	370	Bauhinia .....	364
AMARANTACEÆ .....	360	Bidens .....	394
Amarantus .....	360	Bignonia .....	390
AMARYLLIDACEÆ.....	357	BIGNONIACEÆ.....	390
<i>Amellus</i> .....	395	BIXACEÆ .....	380
Amerimnon .....	363	Bœrhaavia .....	361
Amoreuxia .....	380	BOMBACACEÆ .....	379
ANACARDIACEÆ.....	375	BORRAGINACEÆ.....	385
Anacardium .....	375	Bouteloua .....	350
Andropogon .....	350	<i>Bouteloua</i> .....	350
<i>Andropogon</i> .....	354	Bradburya .....	364
Ancimia .....	348	Bravaisia.....	391
Angelonia .....	389	Brickellia .....	394
Anoda .....	377	Bromelia .....	355
Anona.....	361	BROMELIACEÆ .....	355
ANONACEÆ .....	361	<i>Bromus</i> .....	352
Anthephora .....	350	<i>Brosimum</i> .....	358
Antigonon.....	359	Bryophyllum .....	362
APOCYNACEÆ .....	382	Buceras .....	381
ARACEÆ .....	355	Buchea .....	391
Arachis .....	363	Buchnera.....	389
Argemone.....	362	Buettneria .....	379

	PAGE.		PAGE.
Bumelia.....	376, 404	Citharexylum.....	386
Bunchosia.....	369	Cleome.....	362
BURSERACEÆ.....	370	Clerodendron.....	386
Caesalpinia.....	364	Clitoria.....	366
Cakile.....	362	Clusia.....	380
Caladium.....	355	Cocos.....	355
Calea.....	394	Codiaeum.....	371
Calliandra.....	364	Coffea.....	392
Callisia.....	356	Coix.....	351
Canavalia.....	365	Colubrina.....	376
Canna.....	358	COMBRETACEÆ.....	381
CANNACEÆ.....	358	Commelina.....	356
CAPPARIDACEÆ.....	362, 399	COMMELINACEÆ.....	356
Capparis.....	362	COMPOSITÆ.....	394
Capraria.....	389	Conobea.....	389
CAPRIFOLIACEÆ.....	393	Conocarpus.....	381
Capsicum.....	388	CONVOLVULACEÆ.....	384
Cardiospermum.....	402	Corallocarpus.....	393
Carica.....	381	Corchorus.....	376
CARICACEÆ.....	381	Cordia.....	385
Casimiroa.....	401	Cosmos.....	394
Cassia.....	365	Cracca.....	366
Cassytha.....	362	CRASSULACEÆ.....	362
Castalia.....	361	Cratæva.....	362
Castilloa.....	358	Crescentia.....	390
Casuarina.....	358	Crotalaria.....	366
CASUARINACEÆ.....	358	Croton.....	371
Cayaponia.....	393	CRUCIFERÆ.....	362
Cecropia.....	358	Crusea.....	392
Cedrela.....	370	Cucumis.....	393
Cedronella.....	387	CUCURBITACEÆ.....	393
Ceiba.....	379	CYCADACEÆ.....	350
CELASTRACEÆ.....	375, 401	Cycas.....	350
<i>Celtis</i> .....	399	Cynodon.....	351
Cenchrus.....	351	<i>Cynosurus</i> .....	351
Cestrum.....	388	CYPERACEÆ.....	354
Chamædorea.....	355	Cyperus.....	354
<i>Chamissoa</i> .....	360	Dactyloctenium.....	351
Chara.....	347	<i>Dalbergia</i> .....	363
CHARACEÆ.....	347	Dalea.....	366
Cheilanthes.....	349	<i>Dalea</i> .....	368
CHENOPODIACEÆ.....	359	Dalechampia.....	371
Chenopodium.....	359	Datura.....	388
Chiococca.....	392	Desmanthus.....	367
Chloris.....	351	Diapedium.....	391
Chlorophora.....	358	Dioscorea.....	357
Chrysophyllum.....	382	DIOSCOREACEÆ.....	357
Cissampelos.....	362	Diospyros.....	382

	PAGE.
Dipholis.....	382
Diphysa.....	367
Distichlis.....	351
Dorstenia.....	358
Duranta.....	386
Dysodia.....	395
EBENACEÆ.....	382
Echites.....	382
Eclipta.....	395
Ehretia.....	386
Eleusine.....	351
Eragrostis.....	351
Erigeron.....	395
Ernodea.....	392
Erythrina.....	367
Esenbeckia.....	401
Eupatorium.....	395
Euphorbia.....	371
EUPHORBIACEÆ.....	370, 401
Evolvulus.....	384
Exostemma.....	392
Ficus.....	358
FILICINÆ.....	348
FLACOURTIACEÆ.....	380
Flaveria.....	395
Forchhammeria.....	399
FUNGI.....	347
Galactia.....	367
Gaudichaudia.....	369
Gomphrena.....	360
GOODENIACEÆ.....	394
Gossypium.....	377
Gouania.....	376
Gouinia.....	352
<i>Gouinia</i> .....	352
GRAMINEÆ.....	350
Gronovia.....	381
Guadua.....	352
Guazuma.....	379
GUTTIFERÆ.....	380
Gymnanthes.....	374
<i>Gynoxis</i> .....	396
Gyrocarpus.....	381
Hamelia.....	392
Helianthus.....	395

	PAGE.
Helicteres.....	379
Heliotropium.....	386
Hemionitis.....	349
Henrya.....	391
<i>Herpestis</i> .....	390
Heteropterys.....	369
Hibiscus.....	378
Hippocratea.....	375, 402
HIPPOCRATEACEÆ.....	375, 402
Hura.....	374
Hybanthus.....	404
Hydrocotyle.....	381
Hyptis.....	387
Ichnanthus.....	352
Indigofera.....	367
Ipomœa.....	384
Iresine.....	360
Isocarpha.....	395
Jacquemontia.....	385
Jacquinia.....	382
Jatropha.....	374
Karatas.....	356
Karwinskia.....	376
Kokera.....	360
LABIATÆ.....	387
Lantana.....	386
LAURACEÆ.....	362
LEGUMINOSÆ.....	363
LEMNACEÆ.....	355
Lennoa.....	382
LENNOACEÆ.....	382
Lentinus.....	347
Leonurus.....	387
Leptochloa.....	352
Leucobryum.....	347
LICHENES.....	347
LILIACEÆ.....	357
Lippia.....	387
LOASACEÆ.....	381
Lonicera.....	393
Lueha.....	376
Luffa.....	393
LYTHRACEÆ.....	381

	PAGE.		PAGE.
<i>Maclura</i> .....	358	Nostoc.....	347
Macrosiphonia.....	383	NYCTAGINACEÆ.....	361
Malachra.....	378	NYMPHÆACEÆ.....	361
Malpighia.....	400	Ocimum.....	387
MALPIGHIACEÆ.....	369, 400	Oplismenus.....	352
MALVACEÆ.....	377	<i>Oplismenus</i> .....	353
Malvastrum.....	378	<i>Orthopogon</i> .....	352
Malvaviscus.....	378	OXALIDACEÆ.....	369
Mangifera.....	375	Oxalis.....	369
Manihot.....	374	Pachira.....	379
Martynia.....	391	PALMÆ.....	354
Maytenus.....	375	Panicum.....	353
Maximiliana.....	380	PAPAVERACEÆ.....	362
Meibomia.....	367	Parmentiera.....	390
Melampodium.....	395	Parosela.....	368
Melanthera.....	395	Paspalum.....	353
MELIACEÆ.....	370	Passiflora.....	380
<i>Melicocca</i> .....	403	PASSIFLORACEÆ.....	380
Melochia.....	379	<i>Pauletia</i> .....	364
Melothria.....	393	Paullinia.....	376, 402
MENISPERMACEÆ.....	362	Pectis.....	396
Mertensia.....	399	PEDALIACEÆ.....	391
Metastelma.....	384	Pedilanthus.....	375
Mikania.....	395	Persea.....	362
Milleria.....	395	Petiveria.....	360
Mimosa.....	368	Petrea.....	387
Mirabilis.....	361	Phaseolus.....	368
Momordica.....	394	Phegopteris.....	349
Monniera.....	390	Philodendron.....	355
Montanoa.....	395	Philoxerus.....	360
MORACEÆ.....	358	Phœnix.....	355
Morinda.....	392	Physalis.....	388
Moringa.....	362	PHYTOLACCACEÆ.....	360
MORINGACEÆ.....	362	Pimenta.....	381
Morus.....	359	Piper.....	358
<i>Morus</i> .....	358	PIPERACEÆ.....	358
Mucuna.....	368	Piscidia.....	368
Muntingia.....	377	Pisonia.....	361
Musa.....	358	Pithecoctenium.....	391
MUSACEÆ.....	358	Pittiera.....	393
MUSCI.....	347	PLANTAGINACEÆ.....	392
Myginda.....	401	Plantago.....	392
Myroxylon.....	368	Pluchea.....	396
MYRSINACEÆ.....	382	PLUMBAGINACEÆ.....	382
MYRTACEÆ.....	381	Plumbago.....	382
<i>Myrtus</i> .....	381	Plumeria.....	383
Nerium.....	383	Podopterus.....	359
Nicotiana.....	388		



	PAGE.
POLYGONACEÆ .....	359
Polypodium.....	349
Porphyllum.....	396
Portulaca.....	361
PORTULACACEÆ.....	361
Priva.....	387
Prosopis.....	368
Protium.....	370
Psidium.....	381
Psychotria.....	392
Pterocarpus.....	368
Punica.....	381
Quararibea.....	379
Ramalina.....	347
Randia.....	392
Raphanus.....	362
Rauwolfia.....	383
RHAMNACEÆ.....	376
Rhizophora.....	381
RHIZOPHORACEÆ.....	381
Rhœo.....	356
Rhynchosia.....	368
Rhynchospora.....	354
Ricinus.....	375
Rosmarinus.....	387
RUBIACEÆ.....	392
Ruellia.....	391
RUTACEÆ.....	379, 401
Sabal.....	355
Salicornia.....	360
Salvia.....	387
Salvinia.....	349
Sambucus.....	393
Samyda.....	380
Sanvitalia.....	396
SAPINDACEÆ.....	376, 402
Sapindus.....	402
SAPOTACEÆ.....	382, 404
Scævola.....	394
<i>Schmidelia</i> .....	402
Schwenkia.....	389
Sclerocarpus.....	396
SCROPHULARIACEÆ.....	389
Selaginella.....	349
Senecio.....	396
Serjania.....	376, 403

	PAGE.
Sesbania.....	368
Sesuvium.....	361
Setariopsis.....	354
<i>Sicydium</i> .....	394
Sida.....	378
Simaruba.....	370
SIMARUBACEÆ.....	370
<i>Sinebra</i> .....	350
Siphonoglossa.....	392
Smilax.....	357
SOLANACEÆ.....	388
Solanum.....	388
Spermacoce.....	393
Spironema.....	356
Spondias.....	375
Stemmadenia.....	383
Sterculia.....	379
STERCULIACEÆ.....	379
Stereophyllum.....	348
Stigmaphyllon.....	369
Streptachne.....	354
Stylosanthes.....	369
Suæda.....	360
Suriana.....	370
Synedrella.....	397
Tabernæmontana.....	383
Tabebuia.....	391
Tagetes.....	397
Talisia.....	403
Tamarindus.....	369
Tecoma.....	391
<i>Terminalia</i> .....	381
<i>Tetramerium</i> .....	391
Tetrapteris.....	370
Thevetia.....	383
Thouinia.....	403
Thrinax.....	355
TILIACEÆ.....	376
Tillandsia.....	356
Tithonia.....	397
Tournefortia.....	386
Tragia.....	375
<i>Triæna</i> .....	350
Tribulus.....	370
Triceratia.....	394
Trichilia.....	370
<i>Tricuspis</i> .....	352
Tridax.....	397

	PAGE.		PAGE.
Triumfetta.....	380	Vinca.....	384
Trixis.....	397	VIOLACEÆ.....	404
Tubiflora.....	392	Waltheria.....	380
Turnera.....	380	Wissadula.....	378
TURNERACEÆ.....	380	Wolffia.....	355
ULMACEÆ.....	399	Xanthium.....	397
UMBELLIFERÆ.....	381	Yucca.....	357
Urera.....	359	Zanthoxylum.....	370
URTICACEÆ.....	359	Zebrina.....	356
Urvillea.....	403	Zephyranthes.....	357
Valerianoides.....	392	Zexmenia.....	397
VERBENACEÆ.....	386	ZYGOPHYLLACEÆ.....	370
Viguiera.....	397		

FIELD COLUMBIAN MUSEUM

PUBLICATION 39.

BOTANICAL SERIES.

VOL. I, No. 5.

STUDIES IN THE HERBARIUM.

I.

HIGINBOTHAMIA,

A NEW GENUS, AND OTHER NEW  
DIOSCOREACEÆ.

NEW AMARANTHACEÆ.

BY

EDWIN B. ULINE, PH.D.

CHARLES FREDERICK MILLSPAUGH, M.D.,  
Curator, Department of Botany.



CHICAGO, U. S. A.

August, 1899.

